

- 1 EB 113 B 111 B 113 I B 113

(43) International Publication Date 28 June 2001 (28.06.2001)

PCT

(10) International Publication Number WO 01/46261 A1

- (51) International Patent Classification⁷: C07K 16/24, 16/28, 14/54, A61K 39/395, C07K 19/00, A61P 17/06, 11/06 // C07K 14/715
- (21) International Application Number: PCT/US00/35305
- (22) International Filing Date:

22 December 2000 (22.12.2000)

(25) Filing Language:

English

(26) Publication Language:

English


(30) Priority Data:

09/470,898 60/213,341 23 December 1999 (23.12.1999) US 22 June 2000 (22.06.2000) US

- (71) Applicant: ZYMOGENETICS, INC. [US/US]; 1201 Eastlake Avenue East, Seattle, WA 98102 (US).
- (72) Inventors: THOMPSON, Penny; 9308 151st Avenue SE, Snohomish, WA 98290 (US). FOSTER, Donald, C.; 3002 NE 181st Street, Lake Forest Park, WA 98155 (US). WENFENG, Xu; 12432 54th Avenue W, Mukilteo, WA 98275 (US). MADDEN, Karen, L.; 11148 SE 29th Place, Bellevue, WA 98004 (US). KELLY, James, D.; 3330 W. Mercer Way, Mercer Island, WA 98040 (US). SPRECHER, Cindy, A.; 8207 39th Avenue NE, Seattle, WA 98115 (US). BLUMBERG, Hal; 4620 Sunnyside Avenue N., Seattle, WA 98103 (US). EAGAN, Maribeth, A.; 620 NW 84th Street, Seattle, WA 98117 (US). JASPERS, Stephen, R.; 5829 150th Place S.W., Edmonds, WA 98026 (US). CHANDRASEKHER, Jasmin, A.; 5912 83rd Place SE, Mercer Island, WA 98040 (US). NOVAK, Julia, E.; 10699 Battle Point Drive NE, Bainbridge Island, WA 98110 (US).
- (74) Agent: LUNN, Paul, G.; ZymoGenetics, Inc., 1201 East-lake Avenue East, Seattle, WA 98102 (US).

[Continued on next page]

(54) Title: METHOD FOR TREATING INFLAMMATION



(57) Abstract: A method for treating IL-20 induced inflammation. An antagonist to IL-20 is administered to treat inflammation and associated diseases. The antagonist can be an antibody that binds to IL-20 or its receptor or a soluble receptor that binds to IL-20. Examples of such diseases are adult respiratory disease, psoriasis, eczema, contact dermatitis, atopic dermatitis, septic shock, multiple organ failure, inflammatory lung injury, bacterial pneumonia, inflammatory bowel disease, rheumatoid arthritis, asthma, ulcerative colitis and Crohn's disease.

WO 01/46261 A1

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,

IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

METHOD FOR TREATING INFLAMMATION

BACKGROUND OF THE INVENTION

5

15

20

25

30

The teachings of all of the references cited herein are incorporated in their entirety herein by reference.

Inflammation normally is a localized, protective response to trauma or microbial invasion that destroys, dilutes, or walls-off the injurious agent and the injured tissue. It is characterized in the acute form by the classic signs of pain, heat, redness, swelling, and loss of function. Microscopically, it involves a complex series of events, including dilation of arterioles, capillaries, and venules, with increased permeability and blood flow, exudation of fluids, including plasma proteins, and leukocyte migration into the area of inflammation.

Diseases characterized by inflammation are significant causes of morbidity and mortality in humans. Commonly, inflammation occurs as a defensive response to invasion of the host by foreign, particularly microbial, material. Responses to mechanical trauma, toxins, and neoplasia also may results in inflammatory reactions. The accumulation and subsequent activation of leukocytes are central events in the pathogenesis of most forms of inflammation. Deficiencies of inflammation compromise the host. Excessive inflammation caused by abnormal recognition of host tissue as foreign or prolongation of the inflammatory process may lead to inflammatory diseases as diverse as diabetes, arteriosclerosis, cataracts, reperfusion injury, and cancer, to post-infectious syndromes such as in infectious meningitis, rheumatic fever, and to rheumatic diseases such as systemic lupus erythematosus and rheumatoid arthritis. The centrality of the inflammatory response in these varied disease processes makes its regulation a major element in the prevention control or cure of human disease.

PCT/US00/35305

An important cytokine in the inflammatory process is interleukin-8 (IL-

2

8). IL-8 is a chemokine. It was identified as an agonist for neutrophils on the basis of two effects, chemotaxis and the release of granule enzymes. IL-8 binds to two receptors on neutrophils. IL-8 receptors are also found on monocytes, basophils, and eosinophils. In human fibroblasts, cytomegalovirus has been shown to induce the expression of IL-8 receptors and to replicate more rapidly when cells are exposed to IL-8. IL-8 is a potent chemoattractant for neutrophils; and the early stages of periodontal disease are characterized by the influx of neutrophils. IL-8 is a potent inducer of angiogenesis in several angiogenesis-dependent chronic inflammatory conditions, including rheumatoid arthritis, psoriasis, and idiopathic pulmonary fibrosis. Additionally, IL-8 is an important source of angiogenic activity in human lung cancer. Also, IL-8 expression correlates with experimental metastatic activity of some melanoma cell lines. Thus, an effective method to treat inflammatory diseases would be to administer an agent that would

15

inhibit IL-8.

10

5

BRIEF DESCRIPTION OF THE DRAWINGS

FIGURES: 1-8 are schematic representations of a representative number of embodiments of the IL-20 soluble receptor.

20

25

30

DESCRIPTION OF THE INVENTION

The present invention fills this need by providing for a method for treating inflammation in which IL-20 plays a role, comprising administering to a mammal in need of treatment of inflammation an antagonist to IL-20 or an antagonist to the IL-20 receptor. The antagonist to IL-20 can be an antibody, antibody fragment or single-chain antibody that binds to IL-20, a soluble receptor that binds to IL-20. The antagonist to the IL-20 receptor can be an antibody, antibody fragment, single-chain antibody or small molecule that binds to the IL-20 receptor. Also an anti-sense nucleotide that binds to the mRNA that encodes IL-20 can be used as an antagonist. We have shown that IL-20 up-regulates IL-8. Inflammatory diseases in which IL-8 plays a

significant role, and for which a decrease in IL-8 would be beneficial are, adult respiratory disease (ARD), septic shock, multiple organ failure, inflammatory lung injury such as asthma or bronchitis, bacterial pneumonia, psoriasis and inflammatory bowel disease such as ulcerative colitis and Crohn's disease, eczema, atopic dermatitis and contact dermatitis. Thus, antagonists to IL-20 can be used to treat these diseases.

The teachings of all the references cited herein are incorporated in their entirety by reference.

Definitions

5

20

25

30

Prior to setting forth the invention in detail, it may be helpful to the understanding thereof to define the following terms.

The terms "amino-terminal" and "carboxyl-terminal" are used herein to denote positions within polypeptides. Where the context allows, these terms are used with reference to a particular sequence or portion of a polypeptide to denote proximity or relative position. For example, a certain sequence positioned carboxyl-terminal to a reference sequence within a polypeptide is located proximal to the carboxyl terminus of the reference sequence, but is not necessarily at the carboxyl terminus of the complete polypeptide.

The term "complement/anti-complement pair" denotes non-identical moieties that form a non-covalently associated, stable pair under appropriate conditions. For instance, biotin and avidin (or streptavidin) are prototypical members of a complement/anti-complement pair. Other exemplary complement/anti-complement pairs include receptor/ligand pairs, antibody/antigen (or hapten or epitope) pairs, sense/antisense polynucleotide pairs, and the like. Where subsequent dissociation of the complement/anti-complement pair is desirable, the complement/anti-complement pair preferably has a binding affinity of <10⁹ M⁻¹.

The term "complements of a polynucleotide molecule" is a polynucleotide molecule having a complementary base sequence and reverse orientation as compared to a reference sequence. For example, the sequence 5' ATGCACGGG 3' is complementary to 5' CCCGTGCAT 3'.

15

20

25

The term "contig" denotes a polynucleotide that has a contiguous stretch of identical or complementary sequence to another polynucleotide. Contiguous sequences are said to "overlap" a given stretch of polynucleotide sequence either in their entirety or along a partial stretch of the polynucleotide. For example, representative contigs to the polynucleotide sequence 5'-ATGGCTTAGCTT-3' are 5'-TAGCTTgagtct-3' and 3'-gtcgacTACCGA-5'.

The term "degenerate nucleotide sequence" denotes a sequence of nucleotides that includes one or more degenerate codons (as compared to a reference polynucleotide molecule that encodes a polypeptide). Degenerate codons contain different triplets of nucleotides, but encode the same amino acid residue (i.e., GAU and GAC triplets each encode Asp).

The term "expression vector" is used to denote a DNA molecule, linear or circular, that comprises a segment encoding a polypeptide of interest operably linked to additional segments that provide for its transcription. Such additional segments include promoter and terminator sequences, and may also include one or more origins of replication, one or more selectable markers, an enhancer, a polyadenylation signal, etc. Expression vectors are generally derived from plasmid or viral DNA, or may contain elements of both.

The term "isolated", when applied to a polynucleotide, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such isolated molecules are those that are separated from their natural environment and include cDNA and genomic clones. Isolated DNA molecules of the present invention are free of other genes with which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as promoters and terminators. The identification of associated regions will be evident to one of ordinary skill in the art (see for example, Dynan and Tijan, *Nature 316:774-78* (1985).

An "isolated" polypeptide or protein is a polypeptide or protein that is

found in a condition other than its native environment, such as apart from blood and
animal tissue. In a preferred form, the isolated polypeptide is substantially free of other

10

15

20

25

30

polypeptides, particularly other polypeptides of animal origin. It is preferred to provide the polypeptides in a highly purified form, i.e. greater than 95% pure, more preferably greater than 99% pure. When used in this context, the term "isolated" does not exclude the presence of the same polypeptide in alternative physical forms, such as dimers or alternatively glycosylated or derivatized forms.

The term "operably linked", when referring to DNA segments, indicates that the segments are arranged so that they function in concert for their intended purposes, *e.g.*, transcription initiates in the promoter and proceeds through the coding segment to the terminator.

A "polynucleotide" is a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end.

Polynucleotides include RNA and DNA, and may be isolated from natural sources, synthesized *in vitro*, or prepared from a combination of natural and synthetic molecules. Sizes of polynucleotides are expressed as base pairs (abbreviated "bp"), nucleotides ("nt"), or kilobases ("kb"). Where the context allows, the latter two terms may describe polynucleotides that are single-stranded or double-stranded. When the term is applied to double-stranded molecules it is used to denote overall length and will be understood to be equivalent to the term "base pairs". It will be recognized by those skilled in the art that the two strands of a double-stranded polynucleotide may differ slightly in length and that the ends thereof may be staggered as a result of enzymatic cleavage; thus all nucleotides within a double-stranded polynucleotide molecule may not be paired. Such unpaired ends will in general not exceed 20 nucleotides in length.

A "polypeptide" is a polymer of amino acid residues joined by peptide bonds, whether produced naturally or synthetically. Polypeptides of less than about 10 amino acid residues are commonly referred to as "peptides".

The term "promoter" is used herein for its art-recognized meaning to denote a portion of a gene containing DNA sequences that provide for the binding of RNA polymerase and initiation of transcription. Promoter sequences are commonly, but not always, found in the 5' non-coding regions of genes.

A "protein" is a macromolecule comprising one or more polypeptide chains. A protein may also comprise non-peptidic components, such as carbohydrate

6

groups. Carbohydrates and other non-peptidic substituents may be added to a protein by the cell in which the protein is produced, and will vary with the type of cell. Proteins are defined herein in terms of their amino acid backbone structures; substituents such as carbohydrate groups are generally not specified, but may be present nonetheless.

The term "receptor" denotes a cell-associated protein that binds to a bioactive molecule (i.e., a ligand) and mediates the effect of the ligand on the cell. Membrane-bound receptors are characterized by a multi-domain structure comprising an extracellular ligand-binding domain and an intracellular effector domain that is typically involved in signal transduction. Binding of ligand to receptor results in a conformational change in the receptor that causes an interaction between the effector domain and other molecule(s) in the cell. This interaction in turn leads to an alteration in the metabolism of the cell. Metabolic events that are linked to receptor-ligand interactions include gene transcription, phosphorylation, dephosphorylation, increases in cyclic AMP production, mobilization of cellular calcium, mobilization of membrane lipids, cell adhesion, hydrolysis of inositol lipids and hydrolysis of phospholipids. In general, receptors can be membrane bound, cytosolic or nuclear, monomeric (e.g., thyroid stimulating hormone receptor, beta-adrenergic receptor) or multimeric (e.g., PDGF receptor, growth hormone receptor, IL-3 receptor, GM-CSF receptor, G-CSF receptor, erythropoietin receptor and IL-6 receptor).

The term "secretory signal sequence" denotes a DNA sequence that encodes a polypeptide (a "secretory peptide") that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger polypeptide is commonly cleaved to remove the secretory peptide during transit through the secretory pathway.

Molecular weights and lengths of polymers determined by imprecise analytical methods (e.g., gel electrophoresis) will be understood to be approximate values. When such a value is expressed as "about" X or "approximately" X, the stated value of X will be understood to be accurate to $\pm 10\%$.

25

5

10

15

20

10

20

25

30

Introduction

The present invention relates to the treatment of inflammation and inflammatory diseases in which IL-20 plays a role either in initiating or spreading or maintenance by administering to the afflicted individual an antagonist to IL-20. That antagonist can be an antibody to IL-20, a soluble receptor that binds to IL-20, an antisense molecule or an antibody that binds to either the IL-20RA subunit or the IL-20RB subunit of the IL-20 receptor.

IL-20 is defined and methods for producing it and antibodies to IL-20 are contained in International Patent Application No. PCT/US98/25228, publication no. WO 99/27103, published November 25, 1998 and U.S. Patent Application No. 09/313,458 filed May 17, 1999. The polynucleotide and polypeptide of human IL-20 are represented by SEQ ID NOs: 1 - 4, and mouse IL-20 by SEQ ID NOs: 5-9.

The receptor to IL-20 has been discovered and is a heterodimer comprised of the polypeptide termed 'IL-20RA' (formally called Zcytor7) and a polypeptide termed 'IL-20RB'. The IL-20RA polypeptide, nucleic acid that encodes it, antibodies to IL-20RA, and methods for producing it are disclosed in U.S. Patent No. 5,945,511 issued August 31, 1999. SEQ ID NOs: 10 - 12 are the IL-20RA polynucleotides and polypeptides. The extracellular domain of the human IL-20RA is comprised of a polypeptide selected from the group consisting of SEQ ID NOs: 12, 55, 63 and 65, the full-length receptor subunit being comprised of SEQ ID NO: 11. The extracellular domain of mouse IL-20RA is SEQ ID NO: 38, SEQ ID NO: 37 being the entire mouse IL-20RA.

The extracellular domain of IL-20RB (SEQ ID NOs: 13-14, and a variant SEQ ID NOs: 18 and 19) is comprised of a polypeptide selected from the group consisting of SEQ ID NOs: 15, 59, 61, 67, 68 and 69. Preferably, the extracellular domain of the IL-20RA polypeptide and the extracellular domain of the IL-20RB polypeptide are covalently linked together. In a preferred embodiment one extracellular subunit polypeptide has a constant region of a heavy chain of an immunoglobulin fused to its carboxy terminus and the other extracellular subunit has a constant light chain of an immunoglobulin (Ig) fused to its carboxy terminus such that the two polypeptides come together to form a soluble receptor and a disulfide bond is formed between the heavy and the light Ig chains. In another method, a peptide linker could be fused to the two carboxy-termini of the polypeptides to form a covalently bonded soluble receptor.

8

SEQ ID NOs: 22 and 23 are constructs of the extracellular domain of IL-20RA fused to a mutated human immunoglobulin gamma 1 constant region produced according to the procedure set forth in example 5. SEQ ID NO: 62 is the predicted mature sequence without the signal sequence. SEQ ID NOs: 20 and 21 are constructs of the extracellular domain of IL-20RB fused to wild type human immunoglobulin kappa light chain constant region produced according to the procedure of example 5. SEQ ID NO: 60 is the predicted mature sequence without the signal sequence. Figure 1 depicts the heterotetramer produced by example 5.

SEQ ID NOs: 52 and 53 are constructs of the extracellular domain of IL20RA fused to a mutated human immunoglobulin gamma 1 constant region produced according to the procedure set forth in example 12. SEQ ID NO: 54 is the predicted mature sequence without the signal sequence. SEQ ID NOs: 56 and 57 are constructs of the extracellular domain of IL-20RB fused to wild type human immunoglobulin kappa light chain constant region produced according to the procedure of example 12. SEQ ID NO: 58 is the predicted mature sequence without the signal sequence. The resultant heterotetramer is almost identical to that produced by example 5, the primary difference being the absence of a polypeptide linker between the extracellular domains and the beginning of the Ig constant regions, 22 in Figure 1. Hereinafter, the term "extracellular domain of a receptor" means the extracellular domain of the receptor or a portion of the extracellular domain that is necessary for binding to its ligand, in this case the ligand being IL-20.

One can link together the extracellular domains of IL-20RA and IL-20RB in a number of ways such that the resultant soluble receptor can bind to IL-20. Figures 1-8 illustrate a representative number of embodiments of the present invention. Common elements in each of the drawings are given the same number. Figure 1 represents the embodiment of the present invention produced according to example 5 below. The soluble receptor construct, designated 10, is comprised of two IL-20 binding site polypeptide chains designated 12 and 14. Each binding site is comprised of the extracellular domain of IL-20RA, designated 16, and the extracellular domain of IL-20RB designated 18.

25

30

. 35

The extracellular domain, 16, of IL-20RA is linked to the constant heavy one (CH1) domain, 20, of the human immunoglobulin gamma 1 heavy chain constant region via linker 22, which is SEQ ID NO:72. The CH1 domain, 20, is then linked to the CH2 domain, 24, via hinge region 23. The CH2 domain, 24, is linked to the CH3 domain, 26, via hinge region 25.

Comparing the construct of Figure 1 with SEQ ID NO:22, the extracellular domain, 16, of IL-20RA extends from amino acid residues 36, a valine, to

15

20

25

30

35

and including amino acid residue 249, a glutamine of SEQ ID NO:22. Polypeptide linker, 22, extends from amino acid residue 250, a glycine to and including amino acid residue 264, a serine, of SEQ ID NO:22. The CH1 domain, 22 of Figure 1, extends from amino acid residue 265, an alanine, to and including amino acid residue 362, a valine, of SEQ ID NO:22. Hinge region 23 of Figure 1 extends from amino acid residue 363, a glutamic acid to and including amino acid residue 377, a proline, of SEQ ID NO: 22. Chains 12 and 14 are disulfide-bonded together by means of disulfide bonds 28 and 30. The disulfide bonds are formed between the heavy chains by the cysteine residues at positions 373 and 376 of SEQ ID NO: 22 of each of the two heavy chains.

Extracellular domain, 18, of IL-20RB is linked to the constant region of the human kappa light chain (CL), 34 of Figure 1 via polypeptide linker 32, which is the polypeptide SEQ ID NO: 72. The extracellular domain, 18, of IL-20RB extends from amino acid residue 30, a valine, to and including amino acid residue 230, an alanine, of SEQ ID NO: 20. Polypeptide linker, 32, extends from amino acid residue 231, a glycine, to and including amino acid residue 245, a serine, of SEQ ID NO:20. The kappa constant light region, 34, extends from amino acid residue 246, an arginine, to and including the final amino acid residue 352, a cysteine, of SEQ ID NO:20. The cysteine at position 352 of SEQ ID NO: 20 forms a disulfide bond, 36 in Figure 1, with the cysteine at position 367 of SEQ ID NO: 22. The constant light chain 34 is thus linked to the hinge region, 23, by disulfide bond, 36. In this way, the extracellular domain, 16, of IL-20RB is linked to the extracellular domain, 18, of IL-20RB to form a soluble receptor.

If the cysteine residues at positions 373 and 376 of SEQ ID NO:22 were changed to different amino acid residues, the two IL-20 binding polypeptides, 12 and 14, would not be disulfide bonded together and would form a construct shown in Figure 2. having hinge region, 27.

Figure 3 shows a very simple soluble receptor 38 of the present invention wherein extracellular domain, 16, of IL-20RA is connected to the extracellular domain, 18, of IL-20RB by means of a polypeptide linker, 40. The polypeptide linker extends from the amino terminus of extracellular domain, 16, of IL-20RA and is connected to the carboxyl terminus of the extracellular domain, 18, of IL-20RB. The polypeptide linker should be between 100-240 amino acids in length, preferably about 170 amino acid residues in length. A suitable linker would be comprised of glycine and serine residues. A possible linker would be multiple units of SEQ ID NO: 72, preferably about 12.

Figure 4 shows an embodiment that has the extracellular domain, 16, of IL-20RA linked to the extracellular domain, 18, of IL-20RB by means of linker 40, as in Figure 3. While the extracellular domain, 16, of IL-20RA is linked to the CH1 domain, 20, as in Figure 1 by means of polypeptide linker 42, which should be about 30 amino acid residues in length. An ideal linker would be comprised of glycine and serine as in SEQ ID NO: 72, and the hinge sequence, 23 of Figure 1.

Figure 5 shows another possible embodiment of the present invention. In this embodiment, a polypeptide linker 44 of about 15 amino acid residue, e.g. SEQ ID NO: 72, links the carboxyl terminus of the extracellular domain, 18, of IL-20RB with the amino terminus of the extracellular domain, 16, of IL-20RA. A polypeptide linker 46 of about 30 amino acid residues extends from the carboxy terminus of the extracellular domain, 16, of IL-20RA to the CH2 domain. The carboxyl terminus of linker 46 would preferably be comprised of the hinge region extending from amino acid residue 363, a glutamic acid to and including amino acid residue 377, a proline, of SEQ ID NO: 22. Nonetheless, polypeptide linker 46 would ideally have at least one cysteine residue at its carboxyl terminus so a disulfide bond could be formed.

10

15

20

25

30

35

The soluble IL-20 receptor of Figure 6 is identical to that of Figure 1 except for the CH3 domain, 26 of Figure 1, is not present on the embodiment of Figure 6. The CH3 region begins at amino acid residue 488, a glycine, and extends to the last residue 594 of SEQ ID NO: 22.

Figure 7 shows a soluble IL-20 receptor construct that is identical to the construct of Figure 1 except both the CH2, and CH3 domains are absent. The CH2 and CH3 domains run from amino acid residue 378, an alanine, to the end of the polypeptide sequence of SEO ID NO: 22.

Figure 8 shows a construct wherein both IL-20RA, 16, and IL-20RB have a polypeptide linker, 48, fused to their respective carboxyl termini. Each polypeptide linker has two cysteine residues such that when they are expressed the cysteines form two disulfide bonds, 50 and 52. In this case the polypeptide linker is comprised of the hinge region, 23 in Figure 1. The hinge region is comprised of amino acid residues 363, a glutamine, to and including amino acid residue 377 of SEQ ID NO: 22.

In another aspect of the invention, a method is provided for producing a soluble receptor comprised of extracellular domains of IL-20RA and IL-20RB comprising (a) introducing into a host cell a first DNA sequence comprised of a transcriptional promoter operatively linked to a first secretory signal sequence followed downstream by and in proper reading frame the DNA that encodes the extracellular portion of IL-20RA and the DNA that encodes an immunoglobulin light chain constant

11

region;(b) introducing into the host cell a second DNA construct comprised of a transcriptional promoter operatively linked to a second secretory signal followed downstream by and in proper reading frame a DNA sequence that encodes the extracellular portion of IL-20RB and a DNA sequence that encodes an immunoglobulin heavy chain constant region domain selected from the group consisting of C_H1, C_H2, C_H3 and C_H4; (c) growing the host cell in an appropriate growth medium under physiological conditions to allow the secretion of a fusion protein comprised of the extracellular domain of IL-20RA and IL-20RB; and (d) isolating the polypeptide from the host cell. In one embodiment, the second DNA sequence further encodes an immunoglobulin heavy chain hinge region wherein the hinge region is joined to the heavy chain constant region domain. In another embodiment, the second DNA sequence further encodes an immunoglobulin variable region joined upstream of and in proper reading frame with the immunoglobulin heavy chain constant region.

10

35

In an alternative embodiment, a method is provided for producing a soluble receptor comprised of the extracellular domains of IL-20RA and IL-20RB comprising (a) introducing into a host cell a first DNA sequence comprised of a transcriptional promoter operatively linked to a first secretory signal sequence followed downstream by and in proper reading frame the DNA that encodes the extracellular portion of IL-20RB and the DNA that encodes an immunoglobulin light chain constant 20 region; (b) introducing into the host cell a second DNA construct comprised of a transcriptional promoter operatively linked to a second secretory signal followed downstream by and in proper reading frame a DNA sequence that encodes the extracellular portion of IL-20RA and a DNA sequence that encodes an immunoglobulin heavy chain constant region domain selected from the group consisting of C_H1, C_H2, C_H3 and C_H4; (c) growing the host cell in an appropriate 25 growth medium under physiological conditions to allow the secretion of a dimerized heterodimeric fusion protein comprised of the extracellular domain of IL-20RA and IL-20RB; and (d) isolating the dimerized polypeptide from the host cell. In one embodiment, the second DNA sequence further encodes an immunoglobulin heavy chain hinge region wherein the hinge region is joined to the heavy chain constant region 30 domain. In another embodiment, the second DNA sequence further encodes an immunoglobulin variable region joined upstream of and in proper reading frame with the immunoglobulin heavy chain constant region. (See U.S. Patent No. 5,843,725.)

As used herein, the term "antibodies" includes polyclonal antibodies, affinity-purified polyclonal antibodies, monoclonal antibodies, and antigen-binding fragments, such as F(ab')₂ and Fab proteolytic fragments. Genetically engineered intact antibodies or fragments, such as chimeric antibodies, Fv fragments, single chain

PCT/US00/35305

antibodies and the like, as well as synthetic antigen-binding peptides and polypeptides, are also included. Non-human antibodies may be humanized by grafting non-human CDRs onto human framework and constant regions, or by incorporating the entire nonhuman variable domains (optionally "cloaking" them with a human-like surface by 5 replacement of exposed residues, wherein the result is a "veneered" antibody). In some instances, humanized antibodies may retain non-human residues within the human variable region framework domains to enhance proper binding characteristics. Through humanizing antibodies, biological half-life may be increased, and the potential for adverse immune reactions upon administration to humans is reduced. The binding affinity of an antibody can be readily determined by one of ordinary skill in the art, for example, by Scatchard analysis. A variety of assays known to those skilled in the art can be utilized to detect antibodies that bind to protein or peptide. Exemplary assays are described in detail in Antibodies: A Laboratory Manual, Harlow and Lane (Eds.) (Cold Spring Harbor Laboratory Press, 1988). Representative examples of such assays include: concurrent immunoelectrophoresis, radioimmunoassay, radioimmunoprecipitation, enzyme-linked immunosorbent assay (ELISA), dot blot or Western blot assay, inhibition or competition assay, and sandwich assay.

We have shown that IL-20 up-regulates IL-8. Inflammatory diseases in which IL-8 plays a significant role, and for which a decrease in IL-8 would be beneficial are, adult respiratory disease (ARD), septic shock, multiple organ failure, inflammatory lung injury such as asthma or bronchitis, bacterial pneumonia, psoriasis and inflammatory bowel disease such as ulcerative colitis and Crohn's disease, eczema, atopic dermatitis and contact dermatitis. Thus, antagonists to IL-20 can be used to treat these diseases.

25

30

35

20

10

15

Biology of IL-20, Its receptor and Its Role in Psoriasis

Two orphan class II cytokine receptors, both of which are expressed in skin, were identified as IL-20 receptor subunits. Both IL-20 receptor subunits are required for ligand binding, distinguishing their role from that of subunits in the four other known class II cytokine receptors. IL-20RA and IL-20RB are also coexpressed in a number of human tissues besides skin, including ovary, adrenal gland, testis, salivary gland, muscle, lung, kidney, heart and to a lesser degree the small intestine suggesting additional target tissues for IL-20 action. Additionally, we have detected IL-20RA mRNA but not IL-20RB mRNA in several human tissues including stomach, thyroid, pancreas, uterus, brain and prostate suggesting that IL-20RA may partner with other class II receptor subunits. We conclude that the IL-20 heterodimeric receptor is

structurally similar to other class II cytokine receptors and is expressed in skin where we have demonstrated activity of the IL-20 ligand.

Two lines of evidence indicate that a role IL-20 and its receptor are involved in psoriasis. This multigenic skin disease is characterized by increased keratinocyte proliferation, altered keratinocyte differentiation, and infiltration of immune cells into the skin. The first line of evidence for a role of IL-20 in psoriasis is that the observed hyperkeratosis and thickened epidermis in the transgenic mice that resemble human psoriatic abnormalities. Decreased numbers of tonofilaments, thought to be related to defective keratinization, are a striking feature of human psoriasis. Intramitochondrial inclusions have been found in both chemically induced and naturally occurring hyperplastic skin conditions in mice. The cause of the inclusions and their effects on mitochondrial function, if any, are unknown. We conclude that IL-20 transgenic mice exhibit many of the characteristics observed in human psoriasis.

A second line of evidence that implicates the IL-20 receptor in psoriasis is that both IL-20RA and IL-20RB mRNA are markedly upregulated in human psoriatic skin compared to normal skin. Both IL-20 receptor subunits are expressed in keratinocytes throughout the epidermis and are also expressed in a subset of immune and endothelial cells. We propose that increased expression of an activated IL-20 receptor may alter the interactions between endothelial cells, immune cells and keratinocytes, leading to dysregulation of keratinocyte proliferation and differentiation.

A crucial step in understanding the function of a novel cytokine is the identification and characterization of its cognate receptor. We have successfully used a structure-based approach to isolate a novel interleukin that ultimately led to the isolation of its receptor. IL-20 stimulates signal transduction in the human keratinocyte HaCaT cell line, supporting a direct action of this novel ligand in skin. In addition, IL-1β, EGF and TNF-α, proteins known to be active in keratinocytes and to be involved with proliferative and pro-inflammatory signals in skin, enhance the response to IL-20. In both HaCaT and BHK cells expressing the IL-20 receptor, IL-20 signals through STAT3.

30

35

10

15

20

Use of Antagonist to IL-20 to Treat Psoriasis

As indicated in the discussion above and the examples below, IL-20 is involved in the pathology of psoriasis. The present invention is in particular a method for treating psoriasis by administering antagonists to IL-20. The antagonists to IL-20 can either be a soluble receptor that binds to IL-20 or antibodies, single chain antibodies

15

20

25

30

or fragments of antibodies that bind to either IL-20 or the IL-20 receptor. The antagonists will thus prevent activation of the IL-20 receptor.

Psoriasis is one of the most common dermatologic diseases, affecting up to 1 to 2 percent of the world's population. It is a chronic inflammatory skin disorder characterized by erythematous, sharply demarcated papules and rounded plaques, covered by silvery micaceous scale. The skin lesions of psoriasis are variably pruritic. Traumatized areas often develop lesions of psoriasis. Additionally, other external factors may exacerbate psoriasis including infections, stress, and medications, *e.g.* lithium, beta blockers, and anti-malarials.

PCT/US00/35305

The most common variety of psoriasis is called plaque type. Patients with plaque-type psoriasis will have stable, slowly growing plaques, which remain basically unchanged for long periods of time. The most common areas for plaque psoriasis to occur are the elbows knees, gluteal cleft, and the scalp. Involvement tends to be symmetrical. Inverse psoriasis affects the intertriginous regions including the axilla, groin, submammary region, and navel, and it also tends to affect the scalp, palms, and soles. The individual lesions are sharply demarcated plaques but may be moist due to their location. Plaque-type psoriasis generally develops slowly and runs an indolent course. It rarely spontaneously remits.

Eruptive psoriasis (guttate psoriasis) is most common in children and young adults. It develops acutely in individuals without psoriasis or in those with chronic plaque psoriasis. Patients present with many small erythematous, scaling papules, frequently after upper respiratory tract infection with beta-hemolytic streptococci. Patients with psoriasis may also develop pustular lesions. These may be localized to the palms and soles or may be generalized and associated with fever, malaise, diarrhea, and arthralgias...

About half of all patients with psoriasis have fingernail involvement, appearing as punctate pitting, nail thickening or subungual hyperkeratosis. About 5 to 10 percent of patients with psoriasis have associated joint complaints, and these are most often found in patients with fingernail involvement. Although some have the coincident occurrence of classic Although some have the coincident occurrence of classic rheumatoid arthritis, many have joint disease that falls into one of five type

PCT/US00/35305 WO 01/46261

15

associated with psoriasis: (1) disease limited to a single or a few small joints (70 percent of cases); (2) a seronegative rheumatoid arthritis-like disease; (3) involvement of the distal interphalangeal joints; (4) severe destructive arthritis with the development of "arthritis mutilans"; and (5) disease limited to the spine.

Psoriasis can be treated by administering antagonists to IL-20. The preferred antagonists are either a soluble receptor to IL-20 or antibodies, antibody fragments or single chain antibodies that bind to either the IL-20 receptor or to IL-20. The antagonists to IL-20 can be administered alone or in combination with other established therapies such as lubricants, keratolytics, topical corticosteroids, topical vitamin D derivatives, anthralin, systemic antimetabolites such as methotrexate, psoralen-ultraviolet-light therapy (PUVA), etretinate, isotretinoin, cyclosporine, and the topical vitamin D3 derivative calcipotriol. The antagonists, in particularly the soluble receptor or the antibodies that bind to IL-20 or the IL-20 receptor can be administered to individual subcutaneously, intravenously, or transdermally using a cream or transdermal patch that contains the antagonist of IL-20. If administered subcutaneously, the antagonist can be injected into one or more psoriatic plaques. If administered transdermally, the antagonists can be administered directly on the plaques using a cream containing the antagonist to IL-20.

Use of Antagonists to IL-20 to Treat Inflammatory Conditions of the Lung. 20

Antagonists to IL-20 can be administered to a person who has asthma, bronchitis or cystic fibrosis or other inflammatory lung disease to treat the disease. The antagonists can be administered by any suitable method including intravenous, subcutaneous, bronchial lavage, and the use of inhalant containing an antagonist to IL-20.

Administration of Antagonists to IL-20

5

10

15

25

30

The quantities of antagonists to IL-20 necessary for effective therapy will depend upon many different factors, including means of administration, target site, physiological state of the patient, and other medications administered. Thus, treatment dosages should be titrated to optimize safety and efficacy. Typically, dosages used in vitro may provide useful guidance in the amounts useful for in vivo administration of

WO 01/46261

PCT/US00/35305

these reagents. Animal testing of effective doses for treatment of particular disorders will provide further predictive indication of human dosage. Methods for administration include oral, intravenous, peritoneal, intramuscular, transdermal or administration into the lung or trachea in spray form by means or a nebulizer or atomizer. Pharmaceutically acceptable carriers will include water, saline, buffers to name just a few. Dosage ranges would ordinarily be expected from 1µg to 1000µg per kilogram of body weight per day. A dosage for an average adult of the IL-20 soluble receptor would be about 25 mg given twice weekly as a subcutaneous injection. A dosage of an antibody that binds to IL-20 would be about 25 mg given twice weekly. A mixture of antibodies that bind to the IL-20RA and IL-20RB subunits would be about 25 mg given twice weekly for an adult. Injections could be given at the site of psoriatic lesions for the treatment of psoriasis. For subcutaneous or intravenous administration of the antagonist to IL-20, the antibody or soluble receptor can be in phosphate buffered saline. Also in skin diseases such as psoriasis, the antagonist to IL-20 can be administered via an ointment or transdermal patch. The doses by may be higher or lower as can be determined by a medical doctor with ordinary skill in the art. For a complete discussion of drug formulations and dosage ranges see Remington's Pharmaceutical Sciences, 18th Ed., (Mack Publishing Co., Easton, Penn., 1996), and Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 9th Ed. (Pergamon Press 1996).

16

20

10

15

The invention is further illustrated by the following non-limiting examples.

17

Example 1

Up-regulation of IL-8 by IL-20

Methods:

5

10

15

20

Normal Human Epidermal neonatal keratinocytes (NHEK) (from Clonetics) at passage 2 were plated and grown to confluency in 12 well tissue culture plates. KGM (Keratinocyte growth media) was purchased from Clonetics. When cells reached confluency, they were washed with KGM media minus growth factors = KBM (keratinocyte basal media). Cells were serum starved in KBM for 72 hours prior to the addition of test compounds. Thrombin at 1 I.U./mL and trypsin at 25nM were used as positive controls. One mL of media/well was added. KBM only was used as the negative control.

IL-20 was made up in KBM media and added at varying concentrations, from 2.5µg/ml down to 618ng/mL in a first experiment and from 2.5µg/mL down to 3ng/mL in a second experiment.

Cells were incubated at 37° C, 5% CO₂ for 48 hours. Supernatants were removed and frozen at -80° C for several days prior to assaying for IL-8 and GM-CSF levels. Human IL-8 Immunoassay kit # D8050 (RandD Systems, Inc.) and human GM-CSF Immunoassay kit # HSGMO (RandD Systems, Inc.) were used to determine cytokine production following manufacturer's instructions.

Results

The results indicated that the expression of IL-8 and GM-CSF were induced by IL-20.

Example 2 Cloning of IL-20RB

Cloning of IL-20RB coding region

5

10

15

20

25

30

Two PCR primers were designed based on the sequence from International Patent Application No. PCT/US99/03735 (publication no. WO 99/46379) filed on March 8, 1999. SEQ ID NO: 16 contains the ATG (Met1) codon with an EcoRI restriction site, SEQ ID NO: 17 contains the stop codon (TAG) with an XhoI restriction site. The PCR amplification was carried out using a human keratinocyte (HaCaT) cDNA library DNA as a template and SEQ ID NO: 16 and SEQ ID NO: 17 as primers. The PCR reaction was performed as follows: incubation at 94°C for 1 min followed by 30 cycles of 94°C for 30 sec and 68°C for 2 min, after additional 68°C for 4 min, the reaction was stored at 4°C. The PCR products were run on 1% Agarose gel, and a 1 kb DNA band was observed. The PCR products were cut from the gel and the DNA was purified using a QIAquick Gel Extraction Kit (Qiagen). The purified DNA was digested with EcoRI and XhoI, and cloned into a pZP vector that was called pZP7N. A pZP plasmid is a mammalian expression vector containing an expression cassette having the mouse metallothionein-1 promoter, human tPA leader peptide, multiple restriction sites for insertion of coding sequences, a Glu-Glu tag, and a human growth hormone terminator. The plasmid also has an E. coli origin of replication, a mammalian selectable marker expression unit having an SV40 promoter, an enhancer and an origin of replication, as well as a DHFR gene, and the SV40 terminator. Several IL-20RBpZP7N clones were sequenced. They all contain three non-conservative mutations compared with the sequence of IL-20RB in PCT/US99/03735: (sequence IL-20RBpZP7N), 146 Pro (CCC) - Thr (ACC), 148 His (CAT) -- Asp (GAT), and 171 Thr (ACG) - Arg (AGG).

To verify the three substitutions in IL-20RB-pZP7N clone, PCR amplification was carried out using three difference cDNA sources — fetal skin marathon cDNA, HaCaT cDNA library DNA, and prostate smooth muscle cDNA library DNA — as templates. The PCR products were gel purified and sequenced. The

sequence of each of the three PCR products was consistent with that of the IL-20RB-pZP7N clone. IL-20RB is SEQ ID NO: 13 and 14, and the mature extracellular domain is SEQ ID NO: 15.

5

Example 3

Binding of IL-20 to IL-20RB/ IL-20RA Heterodimer

A cell-based binding assay was used to verify IL-20 binds to IL-20RA-IL-20RB heterodimer.

10

15

20

Expression vectors containing known and orphan Class II cytokine receptors (including IL-20RA and IL-20RB) were transiently transfected into COS cells in various combinations, which were then assayed for their ability to bind biotin-labeled IL-20 protein. The results show IL-20RB- IL-20RA heterodimer is a receptor for IL-20. The procedure used is described below.

The COS cell transfection was performed in a 12-well tissue culture plate as follows: 0.5 µg DNA was mixed with medium containing 5 µl lipofectamine in 92 µl serum free Dulbecco's modified Eagle's medium (DMEM) (55 mg sodium pyruvate, 146 mg L-glutamine, 5 mg transferrin, 2.5 mg insulin, 1 µg selenium and 5 mg fetuin in 500 ml DMEM), incubated at room temperature for 30 minutes and then added to 400 µl serum free DMEM media. This 500 µl mixture was then added to 1.5 x 10⁵ COS cells/well and incubated for 5 hours at 37° C. 500 µl 20% fetal bovine serum (FBS) DMEM media was added and incubated overnight.

The assay, a modification of the "secretion trap" (Davis, S., et al., Cell 87: 1161-1169 (1996), was performed as follows: cells were rinsed with PBS/1% bovine serum albumin (BSA) and blocked for 1 hour with TNB (0.1 M Tris-HCl, 0.15 M NaCl and 0.5% Blocking Reagent (NEN Renaissance TSA-Direct Kit Cat# NEL701) in water). This was followed by a one-hour incubation with 3 μg/ml biotinylated IL-20 protein in TNB. Cells were washed with PBS/1% BSA and incubated for another hour with 1:300 diluted streptavidin-HRP (NEN kit) in TNB. Following another wash, cells were fixed for 15 minutes with 1.8% Formaldehyde in phosphate-buffered saline (PBS).

20

Cells were then washed with TNT (0.1 M Tris-HCL, 0.15 M NaCl, and 0.05% Tween-20 in water). Positive binding signals were detected following a five-minute incubation with fluorescein tyramide reagent diluted 1:50 in dilution buffer (NEN kit). Cells were washed with TNT, preserved with Vectashield Mounting Media (Vector Labs) diluted 1:5 in TNT, and visualized using an FITC filter on an inverted fluorescent microscope.

Example 4

Up-regulation of Inflammatory Cytokines by IL-20

Cell Treatment

5

The human keratinocyte cell line, HaCaT was grown at 37°C to several days post-confluence in T-75 tissue culture flasks. At this point, normal growth media (DMEM + 10% FBS) was removed and replaced with serum-free media. Cells were then incubated for two days at 37°C. DMEM was then removed and four flasks of cells per treatment were treated with one of each of the following conditions for four hours at 37°C: recombinant human (rh) IL-1 alpha at 5 ng/mL, rh IL-1 alpha at 20 ng/mL, rh IL-1 alpha at 5 ng/mL + IL-20 at 1μg/mL, IL-20 at 1μg/mL, or rh IL-10 at 10 ng/mL.

RNA Isolation

using a guanidium thiocyanate solution. Total RNA was isolated from the cell lysate by an overnight spin on a cesium chloride gradient. The following day, the RNA pellet was resuspended in a TE/SDS solution and ethanol precipitated. RNA was then quantitated using a spectrophotometer, followed by a DNase treatment as per Section V.B. of Clontech's AtlasTM cDNA Expression Arrays User Manual (version PT3140-1/PR9X390, published 11/5/99). Quality of RNA samples was verified by purity calculations based on spec readings, and by visualization on agarose gel. Genomic contamination of the RNA samples was ruled out by PCR analysis of the beta-actin gene.

15

20

25

Probe Synthesis

Clontech's protocols for polyA+ enrichment, probe synthesis and hybridization to AtlasTM arrays were followed (see above, plus AtlasTM Pure Total RNA Labeling System User Manual, PT3231-1/PR96157, published 6/22/99). Briefly, polyA+ RNA was isolated from 50 mg of total RNA using streptavidin coated magnetic beads (by Clontech, Paolo Alto, CA) and a magnetic particle separator. PolyA+ RNA was then labeled with ^{alpha32}P-dATP via RT-PCR. Clontech CDS primers specific to the 268 genes on the AtlasTM human cytokine/receptor array (Cat. #7744-1) were used in the reaction. Labeled probe was isolated using column chromatography and counted in scintillation fluid.

Array membrane Hybridization

AtlasTM arrays were pre-hybridized with Clontech ExpressHyb plus 100 mg/mL heat denatured salmon sperm DNA for at least thirty minutes at 68°C with continuous agitation. Membranes were then hybridized with 1.9 x 10⁶ CPM/mL (a total of 1.14 x 10⁷ CPM) overnight at 68°C with continuous agitation. The following day, membranes were washed for thirty minutes x 4 in 2X SSC, 1% SDS at 68°C, plus for thirty minutes x 1 in 0.1X SSC, 0.5% SDS at 68°C, followed by one final room temperature wash for five minutes in 2X SSC. Array membranes were then placed in Kodak plastic pouches sealed and exposed to a phosphor imager screen overnight at room temperature. The next day, phosphor screens were scanned on a phosphor imager and analyzed using Clontech's AtlasImageTM 1.0 software.

Results

Genes Up-regulated by IL-20

- 1. Tumor necrosis factor (TNF) was up-regulated 1.9-2.4 fold by IL-20.
- 2. Placental growth factors 1 & 2 (PLGF) were up-regulated 1.9-2.0 fold by IL-20.
- 3. Coagulating factor II receptor was up-regulated 2.0-2.5 fold by IL-20.
- 30 4. Calcitonin receptor was up-regulated 2.2-2.3 fold by IL-20.

20

30

WO 01/46261 PCT/US00/35305

5. TNF-inducible hyaluronate-binding protein TSG-6 was up-regulated 2.1-2.2 fold by IL-20.

22

- 6. Vascular endothelial growth factor (VEGF) receptor-1 precursor, tyrosine-protein kinase receptor (FLT-1) (SFLT) was up-regulated 2.1-2.7 fold by IL-20.
- 7. MRP-8 (calcium binding protein in macrophages MIF- related) was upregulated 2.9-4.1 fold by IL-20.
 - 8. MRP-14 (calcium binding protein in macrophages MIF-related) was upregulated 3.0-3.8 fold by IL-20.
 - 9. Relaxin H2 was up-regulated 3.14 fold by IL-20.
- 10. Transforming growth factor beta (TGFβ) receptor III 300 kDa was up-regulated2.4-3.6 fold by IL-20.

Genes Showing Synergy with IL-20 + IL-1 Treatment

- 1. Bone morphogenic protein 2a was up-regulated 1.8 fold with IL-20 treatment alone, 2.5 fold with IL-1 treatment alone, and 8.2 fold with both IL-20 and IL-1 treatment together.
 - 2. MRP-8 was up-regulated 2.9 fold with IL-20 treatment alone, 10.7 fold with IL-1 treatment alone and 18.0 fold with both IL-20 and IL-1 treatment together.
- 3. Erythroid differentiation protein (EDF) was up-regulated 1.9 fold with IL-20 treatment alone, 9.7 fold with IL-1 treatment alone and 19.0 fold with both IL-20 and IL-1 treatment together.
- 4. MRP-14 (calcium binding protein in macrophages, MIF related) was upregulated 3.0 fold with IL-20 treatment alone, 12.2 fold with IL-1 treatment alone and 20.3 fold with both IL-20 and IL-1 treatment together.
- 5. Heparin-binding EGF-like growth factor was up-regulated 2.0 fold with IL-20 treatment alone, 14 fold with IL-1 treatment alone and 25.0 fold with both IL-20 and IL-1 treatment together.
 - 6. Beta-thromboglobulin-like protein was up-regulated 1.5 fold with IL-20 treatment alone, 15 fold with IL-1 treatment alone and 27 fold with both IL-20 and IL-1 treatment together.

PCT/US00/35305 WO 01/46261

7. Brain-derived neurotrophic factor (BDNF) was up-regulated 1.7 fold with IL-20 treatment alone, 25 fold with IL-1 treatment alone and 48 fold with both IL-20 and IL-1 treatment together.

23

8. Monocyte chemotactic and activating factor MCAF was up-regulated 1.3 fold with IL-20 treatment alone, 32 fold with IL-1 treatment alone and 56 fold with both IL-20 and IL-1 treatment together.

5

10

15

20

25

30

Example 5

IL-20RA/RB receptor-Ig fusion Heterotetramer

The expression vector pEZE3 was used to express the recombinant IL-20 receptor-Ig fusion protein. The plasmid pEZE3 is derived from pDC312. pDC312 was obtained through license from Immunex Corporation. The plasmids pDC312 and pEZE3 contain an EASE segment as described in WO 97/25420. The presence of the EASE segment in an expression vector can improve expression of recombinant proteins two to eight fold in stable cell pools.

The plasmid pEZE3 is a tricistronic expression vector that may be used to express up to three different proteins in mammalian cells, preferably Chinese Hamster Ovary (CHO) cells. The pEZE3 expression unit contains the cytomegalovirus (CMV) enhancer/promoter, the adenovirus tripartite leader sequence, a multiple cloning site for insertion of the coding region for the first recombinant protein, the policyirus type 2 internal ribosome entry site, a second multiple cloning site for insertion of the coding region for the second recombinant protein, an encephalomyocarditis virus internal ribosome entry site, a coding segment for mouse dihydrofolate reductase, and the SV40 transcription terminator. In addition, pEZE3 contains an E. coli origin of replication and the bacterial beta lactamase gene.

The IL-20 receptor-Ig fusion protein is a disulfide linked heterotetramer consisting of two chains of the extracellular domain of the human IL-20RB fused to the wild type human immunoglobulin kappa light chain constant region and two chains of the human IL-20RA protein extracellular domain fused to a mutated human immunoglobulin gamma 1 constant region. The human immunoglobulin gamma 1

constant region contains amino acid substitutions to reduce FcyRI binding and C1q complement fixation.

The human IL-20RB extracellular domain human immunoglobulin kappa light chain constant region fusion construct was generated by overlap PCR. The IL-20RB coding segment consists of amino acids 1 to 230. The template used for the PCR amplification of the IL-20R segment was generated IL-20RB human kappa light chain constant region expression construct as described below in Example 12. Oligonucleotide primers SEQ ID NO: 24 and SEQ ID NO: 25 were used to amplify the IL-20RB segment. The entire wild type human immunoglobulin kappa light chain 10 constant region was used. The template used for the PCR amplification of the wild type human immunoglobulin kappa light chain constant region segment was generated IL-20RB human kappa light chain constant region expression construct as described in Example 12. Oligonucleotide primers SEQ ID NO: 26 and SEQ ID NO: 27 were used to amplify the wild type human immunoglobulin kappa light chain constant region. The two protein coding domains were linked by overlap PCR using oligonucleotides 15 SEQ ID NO: 24 and SEQ ID NO: 27. A (Gly₄Ser)₃ (SEQ ID NO: 72) peptide linker was inserted between the two protein domains. The (Gly₄Ser)₃ peptide linker was encoded on the PCR primers SEQ ID NO: 26 and SEQ ID NO:25. The resultant IL-20RB extracellular domain/kappa light chain constant region fusion construct is shown by SEQ ID NOs: 20 and 21. The predicted mature polypeptide, minus the signal 20 sequence, is SEQ ID NO: 60. The portion of the extracellular domain of IL-20RB that was actually used was comprised of the amino acid sequence of SEQ ID NO: 61. Nterminal sequencing resulted in the predicted amino acid sequence.

The human IL-20RA extracellular domain human immunoglobulin

gamma 1 heavy chain constant region fusion construct was generated by overlap PCR of four separate DNA fragments, each generated by separate PCR amplification reactions. The first fragment contained an optimized tPA (tissue plasminogen activator) signal sequence. The tPA signal sequence was amplified using oligonucleotide primers SEQ ID NO: 28 and SEQ ID NO: 29 using an in-house previously generated expression vector as the template. The second fragment contained the IL-20RA extracellular domain-coding region consisting of amino acids 30 to 243 of SEQ ID NO: 11.

10

15

20

25

30

PCT/US00/35305

Oligonucleotide primers SEQ ID NO: 30 and SEQ ID NO: 31 were used to amplify this IL-20RA segment using a previously generated clone of IL-20RA as the template.

The human gamma 1 heavy chain constant region was generated from 2 segments. The first segment containing the C_H1 domain was amplified using oligonucleotide primers SEQ ID NO: 32 and SEQ ID NO: 33 using a clone of the wild type human gamma 1 heavy chain constant region as the template. The second segment containing the remaining hinge, C_H2, and C_H3 domains of the human immunoglobulin gamma 1 heavy chain constant region was generated by PCR amplification using oligonucleotide primers SEQ ID NO: 34 and SEQ ID NO: 35. The template used for this PCR amplification was from a previously generated human gamma 1 Fc construct that contained codons for amino acid substitutions to reduce FcγRI binding and C1q complement fixation as described in Example 12.

The four protein coding domains were linked by overlap PCR using oligonucleotides SEQ ID NO: 28 and SEQ ID NO: 35. A (Gly₄Ser)₃ peptide linker was inserted between the IL-20RA and CH1 protein domains. The (Gly₄Ser)₃ peptide linker was encoded on the PCR primers SEQ ID NO: 32 and SEQ ID NO: 31. The IL-20RA extracellular domain/ domain human immunoglobulin gamma 1 heavy constant region fusion protein and DNA sequence are shown in SEQ ID NOs: 22 and 23. The predicted mature polypeptide sequence, minus the signal sequence, is SEQ ID NO: 62. The portion of extracellular domain of IL-20RA that was actually used was comprised of SEQ ID NO: 63.

The IL-20RB extracellular domain human immunoglobulin kappa light chain constant region fusion coding segment was cloned into the second MCS while the human IL-20RA extracellular domain human immunoglobulin gamma 1 heavy chain constant region fusion coding segment was cloned into the first MCS of pEZE3. The plasmid was used to transfect CHO cells. The cells were selected in medium without hypoxanthine or thymidine and the transgene was amplified using methotrexate. The presence of protein was assayed by Western blotting using anti human gamma 1 heavy chain constant region and anti human kappa light chain antibodies. N-terminal sequencing revealed that the optimized tPA leader was not completely cleaved. The observed mass indicated that the first residue of the polypeptide sequence to be

pyroglutamic acid, and the N-terminal sequence appears to be pyroEEIHAELRRFRRVPCVSGG (SEQ ID NO: 64), the underlined portion being remnants of the tPA leader.

Example 6

5

10

15

20

25

30

IL-20 Transgenic Phenotype

Both human and mouse IL-20 were overexpressed in transgenic mice using a variety of promoters. The liver-specific mouse albumin promoter, directing expression of human IL-20, was used initially in an attempt to achieve circulating levels of protein. Subsequent studies were conducted using the keratin 14 (K14) promoter, which primarily targets expression to the epidermis and other stratified squamous epithelia; the mouse metallothionein-1 promoter, which gives a broad expression pattern; and the EµLCK promoter, which drives expression in cells of the lymphoid lineage. Similar results were obtained in all four cases, possibly because these promoters all give rise to circulating levels of IL-20.

In all cases, transgenic pups expressing the IL-20 transgene were smaller than non-transgenic littermates, had a shiny appearance with tight, wrinkled skin and died within the first few days after birth. Pups had milk in their stomachs indicating that they were able to suckle. These mice had swollen extremities, tail, nostril and mouth regions and had difficulty moving. In addition, the mice were frail, lacked visible adipose tissue and had delayed ear and toe development. Low expression levels in liver (less than 100 mRNA molecules/cell) were sufficient for both the neonatal lethality and skin abnormalities. Transgenic mice without a visible phenotype either did not express the transgene, did not express it at detectable levels, or were mosaic.

Histologic analysis of the skin of the IL-20 transgenic mice showed a thickened epidermis, hyperkeratosis and a compact stratum corneum compared to non-transgenic littermates. Serocellular crusts (scabs) were observed occasionally. Electron microscopic (EM) analysis of skin from transgenic mice showed intramitochondrial lipoid inclusions, mottled keratohyaline granules, and relatively few tonofilaments similar to that observed in human psoriatic skin and in mouse skin disease models. In addition, many of the transgenic mice had apoptotic thymic lymphocytes. No other abnormalities were detected by histopathological analysis. These histological and EM

10

15

20

25

results support and extend the observed gross skin alterations.

Example 7

Specificity and Affinity of IL-20 for Its Receptor

The specificity and affinity of IL-20 for its receptor was determined using BHK cells stably transfected with IL-20RA, IL-20RB or both receptor subunits. Binding assays using radiolabeled ligand demonstrated that IL-20 bound to BHK transfectants expressing both IL-20RA and IL-20RB but not to untransfected cells nor to transfectants expressing either receptor subunit alone. Binding of ¹²⁵I-labeled IL-20 was eliminated in the presence of 100-fold excess of unlabeled IL-20 but not with 100-fold excess of the unrelated cytokine, IL-21. The binding affinity (k_D) of IL-20 to the IL-20RA/IL-20RB heterodimeric receptor was determined to be approximately 1.5 nM.

Example 8

IL-20 receptor Activation

To determine if IL-20 binding leads to receptor activation, the factor-dependent pre-B cell line BaF3 was co-transfected with IL-20RA and IL-20RB and treated with IL-20 at various concentrations. IL-20 stimulated proliferation in a dose-dependent manner and gave a detectable signal at 1.1 pM, with a half maximal response at 3.4 pM. We note that the IL-20 concentration for the half maximal proliferative response in BaF3 cells is 1000X lower than that for half maximal binding affinity in BHK cells. Possible explanations for this large difference include the use of different cell lines, different receptor expression levels and different assay outputs. IL-20 also stimulated signal transduction in the biologically relevant human keratinocyte cell line HaCaT, which naturally expresses IL-20RA and IL-20RB. Therefore, IL-20 binds and activates the heterodimeric IL-20RA/IL-20RB receptor at concentrations expected for a cytokine. While the negative controls containing untransfected BaF3

WO 01/46261

Example 9

28

Expression Analysis of IL-20RA and IL-20RB

RT-PCR analysis was performed on a variety of human tissues to determine the expression pattern of IL-20RA and IL-20RB. Both receptor subunits are most highly expressed in skin and testis. The significant result is that IL-20RA and IL-20RB are both expressed in skin, where they have been shown to mediate the IL-20induced response. Both IL-20RA and IL-20RB are also both expressed in monocytes, lung, ovary, muscle, testis, adrenal gland, heart, salivary gland and placenta. IL-20RA is also in brain, kidney, liver, colon, small intestine, stomach, thyroid, pancreas, uterus and prostate while IL-20RB is not.

Example 10

10

15

20

25

30

IL-20RA and IL-20RB mRNA are Up-regulated in Psoriasis In situ hybridization was used to determine whether IL-20 receptor expression is altered in psoriasis. Skin samples from four psoriasis patients and three unaffected patients were assayed with probes specific for the two-receptor subunit mRNAs. All four psoriatic skin samples had high levels of IL-20RA and IL-20RB mRNA in keratinocytes whereas normal skin samples did not have detectable levels of either receptor subunit mRNA. Positive signals in psoriatic skin were also observed in mononuclear immune cells and in endothelial cells in a subset of vessels. Therefore, both IL-20RA and IL-20RB are expressed in keratinocytes, immune cells and endothelial cells, the major cell types thought to interact in psoriasis.

Example 11

Cloning of Mouse IL-20RA

A cross-species hybridization probe was generated which contained the full-length cDNA fragment encoding human IL-20RA. A Southern blot of mouse genomic DNA and Northern blots of mouse RNA were performed to demonstrate that the human IL-20RA cDNA could specifically hybridize to mouse sequences. The Northern blot results indicated that mouse IL-20RA RNA was present in mouse embryo day 15 and 17 as well as heart, brain, lung, liver, kidney, testes, spleen, thymus, liver, stomach, and small intestine.

The human IL-20RA full length DNA hybridization probe was used to screen a mouse genomic library. The library, which was obtained from Clontech (Palo Alto, CA), was generated from an MboI partial digest of mouse genomic DNA and cloned into the BamHI site of Lambda bacteriophage EMBL3 SP6/T7. Positive bacteriophage was plaque purified and bacteriophage DNA was prepared using Promega's Wizard Lambda Preps DNA Purification System. Two genomic restriction enzyme fragments, a 5.7 kb EcoRI fragment and an 8.0 kb SacI fragment, were generated from the positive bacteriophage and subcloned into pBluescript. DNA sequence analysis revealed the presence of 3 exons from the mouse ortholog to human IL-20RA.

PCR primers from the 5' UTR, SEQ ID NO: 40, and 3' UTR, SEQ ID NO: 41, were designed to generate a full-length mouse IL-20RA sequence by PCR amplification. Mouse embryo 15day plus 17 day cDNA was used as the template for the PCR amplification. PCR products were subcloned and sequenced for confirmation. The mouse sequences are SEQ ID NOs: 36 and 37. The mature extracellular domain is comprised of SEQ ID NO: 38.

20

25

30

10

15

Example 12

Construction of an IL-20 Receptor Heterotetramer

A vector expressing a secreted hIL-20RA/hIL-20B heterodimer was constructed. In this construct, the extracellular domain of hIL-20RA was fused to the heavy chain of IgG gamma 1 (IgGγ1), while the extracellular portion of IL-20RB was fused to human kappa light chain (human κ light chain).

Construction of IgG gamma 1 and human κ light fusion vectors

The heavy chain of IgGγ1 was cloned into the Zem229R mammalian expression vector (ATCC deposit No. 69447) such that any extracellular portion of a receptor having a 5' EcoRI and 3' NheI site can be cloned in, resulting in an N-terminal extracellular domain-C-terminal IgGγ1 fusion. The IgGγ1 fragment used in this construct was made by using PCR to isolate the IgGγ1 sequence from a Clontech

human fetal liver cDNA library as template. A PCR reaction using oligos SEO ID NO: 42 and SEQ ID NO: 43 was run as follows: 40 cycles of 94° for 60 sec., 53°C for 60 sec., and 72° for 120 sec.; and 72°C for 7 minutes. PCR products were separated by agarose gel electrophoresis and purified using a QiaQuick™ (Qiagen Inc., Valencia, CA) gel extraction kit. The isolated, 990 bp, DNA fragment was digested with MluI and EcoRI (Boerhinger-Mannheim), extracted with QiaQuick™ gel extraction kit and ligated with oligos SEQ ID NO: 44 and SEQ ID NO: 45, which comprise an MluI/EcoRI linker, into Zem229R previously digested with MluI and EcoRI using standard molecular biology techniques disclosed herein. This generic cloning vector 10 was called Vector#76 hlgGgamma1 w/ Ch1 #786 Zem229R (Vector #76). The polynucleotide sequence of the extracellular domain of hIL-20RA fused to the heavy chain of IgG gamma 1 is show in SEQ ID NO: 52 and the corresponding polypeptide sequence shown in SEQ ID NO: 53, the mature polypeptide, minus the signal sequence being comprised of SEQ ID NO: 54. The portion of the extracellular domain of IL-15 20RA used was comprised of SEQ ID NO: 55.

The human κ light chain was cloned in the Zem228R mammalian expression vector (ATCC deposit No. 69446) such that any extracellular portion of a receptor having a 5' EcoRI site and a 3' KpnI site can be cloned in, resulting in an Nterminal extracellular domain-C-terminal human κ light chain fusion. The human κ light chain fragment used in this construct was made by using PCR to isolate the human κ light chain sequence from the same Clontech hFetal Liver cDNA library used above. A PCR reaction was run using oligos SEQ ID NO: 46 and SEQ ID NO: 47. PCR products were separated by agarose gel electrophoresis and purified using a QiaQuick™ (Qiagen) gel extraction kit. The isolated, 315 bp, DNA fragment was digested with MluI and EcoRI (Boerhinger-Mannheim), extracted with QiaQuick™ gel extraction kit and ligated with the Mlul/EcoRI linker described above, into Zem228R previously digested with MluI and EcoRI using standard molecular biology techniques disclosed herein. This generic cloning vector was called Vector #77 hklight #774 Zem228R (Vector #77). The polynucleotide sequence of the extracellular portion of IL-20RB fused to human kappa light chain is shown in SEQ ID NO: 56 and the corresponding polypeptide sequence shown in SEQ ID NO: 57, the mature polypeptide, minus the

20

25

30

WO 01/46261

31

signal sequence, is comprised of SEQ ID NO: 58. The portion of the extracellular domain of IL-20RB actually used was comprised of SEQ ID NO: 59.

5

10

15

20

25

Insertion of hIL-20RA and IL-20RB extracellular domains into fusion vector constructs

Using the construction vectors above, a construct having human IL-20RA fused to IgGγ1 was made. This construction was done by using PCR to obtain human IL-20RA receptor from hIL-20RA /IgG Vector #102 with oligos SEQ ID NO: 48 and SEQ ID NO: 49 under conditions described as follows: 30 cycles of 94°C for 60 sec., 57°C for 60 sec., and 72°C for 120 sec.; and 72°C for 7 min. The resulting PCR product was digested with EcoRI and NheI, gel purified, as described herein, and ligated into a previously EcoRI and NheI digested and band-purified Vector #76 (above). The resulting vector was sequenced to confirm that the human IL-20Rα /IgG gamma 1 fusion (hIL-20RA /Ch1 IgG) was correct. The hIL-20RA /Ch1 IgG gamma 1 #1825 Zem229R vector was called vector #195. The IL-20RA/Ch1 IgGy1 sequence thus obtained is depicted by SEQ ID NOs: 52 and 53. N-terminal sequencing indicated the presence of the predicted mature polypeptide sequence of SEQ ID NO: 54.

A separate construct having IL-20RB fused to k light was also constructed. The IL-20RB/human κ light chain construction was performed as above by PCRing from DR1/7N-4 with oligos SEQ ID NO: 50 and SEQ ID NO: 51, digesting the resulting band with EcoRI and KpnI and then ligating this product into a previously EcoRI and KpnI digested and band-purified Vec#77 (above). The resulting vector was sequenced to confirm that the IL-20RB/ human κ light chain fusion (IL-20RB/κlight) was correct. This IL-20RB//klight construct is shown by SEQ ID NOs: 56 and 57. Nterminal sequencing of the resultant polypeptide indicated the presence of the predicted mature amino acid sequence comprised of SEQ ID NO: 58. SEQ ID NO:59 is the mature portion of the extracellular domain of IL-20RB used.

Co-expression of the human IL-20RA and human IL-20RB receptors

Approximately 16µg of each of vectors #194 and #195, above, were co-30 transfected into BHK-570 cells (ATCC No. CRL-10314) using Lipofectamine™ reagent (Gibco/BRL), as per manufacturer's instructions. The transfected cells were selected

10

15

20

25

30

for 10 days in DMEM + 5%FBS (Gibco/BRL) containing 1μM of methotrexate (MTX) (Sigma, St. Louis, MO) and 0.5mg/ml G418 (Gibco/BRL) for 10 days. The resulting pool of transfectants was selected again in 10μM MTX and 0.5mg/ml G418 for 10 days.

The resulting pool of doubly selected cells was used to generate protein. Three factories (Nunc, Denmark) of this pool were used to generate 8 L of serum free conditioned medium. This conditioned media was passed over a 1 ml protein-A column and eluted in (10) 750 microliter fractions. 4 of these fractions found to have the highest concentration were pooled and dialyzed (10 kD MW cutoff) against PBS. Finally, the dialyzed material was analyzed by BCA (Pierce) and found to have a concentration of 317 μ g/ml. A total of 951 μ g was obtained from this 8 L purification.

Example 13

IL-20 Binding Activates STAT3 in the HaCaT Keratinocyte Cell Line

IL-20 binds cell lines transfected with both subunits of its receptor. However, these cell lines overexpress the IL-20 receptor relative to its normal level and their relevance to the physiological role of IL-20 is unclear. The human HaCaT keratinocyte cell line, which expresses endogenous IL-20RA and IL-20RB was used to examine IL-20 signal transduction in a biologically relevant cell type. HaCaT cells were infected with recombinant adenovirus containing a reporter construct to allow detection of intracellular signaling. The construct consists of the firefly luciferase gene driven by promoter/enhancer sequences comprised of the serum response element (SRE) and signal transducers and activators of transduction elements (STATs). This assay system detects productive ligand-receptor interactions and indicates possible downstream signal transduction components involved in receptor activation. Treatment with IL-20 alone resulted in a dose-dependent increase in luciferase activity with a half maximal response occurring at approximately 2.3 nM. Subsequent luciferase reporter assays using adenovirus vectors containing only the SRE element or only the STAT elements produced detectable reporter activation only through STATs.

To determine if other cytokines act in concert with IL-20, HaCaT cells were treated with IL-20 alone or in combination with a single submaximal dose of EGF.

IL-1β, or TNFα. In the presence of each of these three proteins, IL-20 treatment resulted in a dose-dependent increase in luciferase activity. IL-20 in combination with IL-1β results in a half-maximal response at approximately 0.5 nM, about five-fold lower than with IL-20 alone. In addition, activation of the reporter gene is detectable at 0.1 nM IL-20, a dose that is at least tenfold lower than the IL-20 dose required alone.

5

10

15

20

25

30

BHK cells transfected with IL-20RA, IL-20RB or both receptor subunits were used to determine whether receptor pairing was required for IL-20 stimulation of STAT-luciferase. As was the case with binding assays, only cells transfected with both receptor subunits responded to IL-20 and did so with a half-maximal response of 5.7 pM. We note that the IL-20 concentration for the half-maximal response in BHK cells is 400-fold lower than that for half-maximal response in HaCaT cells. It is likely that a lower concentration of IL-20 is needed for half-maximal response in BHK cells, as compared to HaCaT cells, due to higher receptor levels in the BHK IL-20 receptor transfectants.

A nuclear translocation assay was used to identify STAT proteins involved in IL-20 action. Both HaCaT cells, with endogenous IL-20 receptors, and BHK cells transfected with IL-20RA and IL-20RB, were treated with IL-20 protein and translocation of STAT3 and STAT1 transcription factors from the cytoplasm to the nucleus was assayed by immunofluorescence.

In unstimulated HaCaT cells, STAT3 staining was predominantly in the cytosol. Treatment of HaCaT cells with IL-20 resulted in a distinct accumulation of STAT3 in the nucleus. Nuclear translocation of STAT3 in response to increasing concentrations of IL-20 occurred with a half-maximal IL-20 concentration of 7 nM. In contrast to STAT3 translocation, HaCaT cells treated with IL-20 did not show any detectable nuclear accumulation of STAT1.

BHK cells transfected with IL-20RA and IL-20RB were used to confirm that the IL-20 receptor was required for IL-20 stimulation of STAT3 nuclear translocation. In BHK cells lacking the IL-20 receptor, STAT3 remained cytosolic following treatment with IL-20. In contrast, in BHK cells transfected with the IL-20 receptor, STAT3 translocated to the nucleus in response to IL-20. Again, STAT1

34

remained cytosolic regardless of IL-20 treatment or IL-20 receptor expression. We conclude that the IL-20 receptor is required for IL-20-mediated STAT3 activation.

5

WHAT IS CLAIMED IS:

1. A method for treating a mammal afflicted with a disease in which an IL-20 polypeptide plays a role, wherein the IL-20 polypeptide is comprised of an amino acid sequence selected from the group consisting of SEQ ID NOs: 1, 2, 3, 4, 5, 6, 7, 8 and 9, comprising:

administering antagonist of said IL-20 polypeptide to said individual.

2. The method of claim 1 wherein the antagonist is selected from the group consisting of an antibody, antibody fragment or single chain antibody that binds to said IL-20 polypeptide, a soluble receptor that binds to said IL-20 polypeptide and an antibody, antibody fragment or single chain antibody which binds to the soluble receptor of IL-20 wherein said soluble receptor is comprised of an IL-20RA subunit and an IL-20RB subunit.

- 3. The method of claim 2 wherein the soluble receptor is a comprised of the extracellular domain of IL-20RA and the extracellular domain of IL-20RB.
- 4. The method of claim 3 wherein the extracellular domain of IL-20RA is fused to a constant region heavy chain of an immunoglobulin (Ig) molecule, and the extracellular domain of IL-20RB is fused to a constant region of a light chain of an Ig molecule.
- 5. The method of claim 2 wherein the antibody, antibody fragment or single chain antibody, which binds to the IL-20 receptor, binds to the IL-20RA subunit.
 - 6. The method of claim 2 wherein the antibody, antibody fragment or single chain antibody, which binds to the IL-20 receptor, binds to the IL-20RB subunit.

WO 01/46261 PCT/US00/35305

- 7. The method of claim 1 wherein the disease that the IL-20 polypeptide plays a role is a skin disease selected from the group consisting of psoriasis, eczema, atopic dermatitis and contact dermatitis.
- 8. The method of claim 1 wherein the disease that the IL-20 polypeptide plays a role is an inflammatory lung selected from the group consisting of adult respiratory disease, asthma, bronchitis and pneumonia.
- 9. A method for promoting the expression of IL-8 in a cell comprising bringing the cell into contact with IL-20.
 - 10. A method for increasing the expression of IL-8 in an individual comprising administering IL-20 to said individual.




FIG.1 SUBSTITUTE SHEET (RULE 26)




FIG.2 SUBSTITUTE SHEET (RULE 26)




FIG.3 SUBSTITUTE SHEET (RULE 26)




FIG.4 SUBSTITUTE SHEET (RULE 26)




FIG.5 SUBSTITUTE SHEET (RULE 26)

WO 01/46261 PCT/US00/35305




FIG.6 SUBSTITUTE SHEET (RULE 26)




FIG.7 SUBSTITUTE SHEET (RULE 26)




FIG.8 SUBSTITUTE SHEET (RULE 26)

SEQUENCE LISTING

<110> ZymoGenetics. Inc. <120> Method for Treating Inflammation <130> 99-108PC <150> 09/470,898 <151> 1999-12-23 <150> 60/213,341 <151> 2000-06-22 <160> 72 <170> FastSEQ for Windows Version 3.0 <210> 1 <211> 176 <212> PRT <213> Homo sapiens <400> 1 Met Lys Ala Ser Ser Leu Ala Phe Ser Leu Leu Ser Ala Ala Phe Tyr Leu Leu Trp Thr Pro Ser Thr Gly Leu Lys Thr Leu Asn Leu Gly Ser Cys Val Ile Ala Thr Asn Leu Gln Glu Ile Arg Asn Gly Phe Ser Asp Ile Arg Gly Ser Val Gln Ala Lys Asp Gly Asn Ile Asp Ile Arg Ile 55 Leu Arg Arg Thr Glu Ser Leu Gln Asp Thr Lys Pro Ala Asn Arg Cys 70 75 Cys Leu Leu Arg His Leu Leu Arg Leu Tyr Leu Asp Arg Val Phe Lys Asn Tyr Gln Thr Pro Asp His Tyr Thr Leu Arg Lys Ile Ser Ser Leu 100 105 110 Ala Asn Ser Phe Leu Thr Ile Lys Lys Asp Leu Arg Leu Cys His Ala 120 125 His Met Thr Cys His Cys Gly Glu Glu Ala Met Lys Lys Tyr Ser Gln 130 135 140

```
Ile Leu Ser His Phe Glu Lys Leu Glu Pro Gln Ala Ala Val Val Lys
                    150
                                        155
Ala Leu Gly Glu Leu Asp Ile Leu Leu Gln Trp Met Glu Glu Thr Glu
                                    170
                165
                                                        175
     <210> 2
     <211> 152
     <212> PRT
     <213> Homo sapiens
     <400> 2
Leu Lys Thr Leu Asn Leu Gly Ser Cys Val Ile Ala Thr Asn Leu Gln
Glu Ile Arg Asn Gly Phe Ser Asp Ile Arg Gly Ser Val Gln Ala Lys
Asp Gly Asn Ile Asp Ile Arg Ile Leu Arg Arg Thr Glu Ser Leu Gln
                            40
Asp Thr Lys Pro Ala Asn Arg Cys Cys Leu Leu Arg His Leu Leu Arg
Leu Tyr Leu Asp Arg Val Phe Lys Asn Tyr Gln Thr Pro Asp His Tyr
                    70
                                        75
Thr Leu Arg Lys Ile Ser Ser Leu Ala Asn Ser Phe Leu Thr Ile Lys
                85
Lys Asp Leu Arg Leu Cys His Ala His Met Thr Cys His Cys Gly Glu
            100
                                105
Glu Ala Met Lys Lys Tyr Ser Gln Ile Leu Ser His Phe Glu Lys Leu
                            120
                                                125
Glu Pro Gln Ala Ala Val Val Lys Ala Leu Gly Glu Leu Asp Ile Leu
    130
                        135
                                            140
Leu Gln Trp Met Glu Glu Thr Glu
145
                    150
      <210> 3
      <211> 151
      <212> PRT
      <213> Homo sapiens
      <400> 3
Met Lys Ala Ser Ser Leu Ala Phe Ser Leu Leu Ser Ala Ala Phe Tyr
```

30

Leu Leu Trp Thr Pro Ser Thr Gly Leu Lys Thr Leu Asn Leu Gly Ser

25

Cys Val Ile Ala Thr Asn Leu Gln Glu Ile Arg Asn Gly Phe Ser Asp Ile Arg Gly Ser Val Gln Ala Lys Asp Gly Asn Ile Asp Ile Arg Ile Leu Arg Arg Thr Glu Ser Leu Gln Asp Thr Lys Pro Ala Asn Arg Cys 70 75 Cys Leu Leu Arg His Leu Leu Arg Leu Tyr Leu Asp Arg Val Phe Lys 85 90 Asn Tyr Gln Thr Pro Asp His Tyr Thr Leu Arg Lys Ile Ser Ser Leu 100 105 110 Ala Asn Ser Phe Leu Thr Ile Lys Lys Asp Leu Arg Leu Cys Leu Glu 120 Pro Gln Ala Ala Val Val Lys Ala Leu Gly Glu Leu Asp Ile Leu Leu 130 135 140 Gln Trp Met Glu Glu Thr Glu 145 150

<210> 4

<211> 127

<212> PRT

<213> Homo sapiens

<400> 4

 Leu Lys Thr Leu Asn Leu Gly Ser Cys Val Ile Ala Thr Asn Leu Gln

 1
 5
 10
 15

 Glu Ile Arg Asn Gly Phe Ser Asp Ile Arg Gly Ser Val Gln Ala Lys 20
 25
 30

 Asp Gly Asn Ile Asp Ile Arg Ile Leu Arg Arg Thr Glu Ser Leu Gln 35
 40
 45

 Asp Thr Lys Pro Ala Asn Arg Cys Cys Leu Leu Arg His Leu Leu Arg 50
 55
 60

 Leu Tyr Leu Asp Arg Val Phe Lys Asn Tyr Gln Thr Pro Asp His Tyr 75
 80

 Thr Leu Arg Lys Ile Ser Ser Leu Ala Asn Ser Phe Leu Thr Ile Lys 90
 95

 Lys Asp Leu Arg Leu Cys Leu Glu Pro Gln Ala Ala Val Val Lys Ala 100
 105
 110

 Leu Gly Glu Leu Asp Ile Leu Leu Gln Trp Met Glu Glu Thr Glu 115
 120
 125

<210> 5

<211> 176

<212> PRT

<213> Mus musculis

<400> 5 Met Lys Gly Phe Gly Leu Ala Phe Gly Leu Phe Ser Ala Val Gly Phe Leu Leu Trp Thr Pro Leu Thr Gly Leu Lys Thr Leu His Leu Gly Ser Cys Val Ile Thr Ala Asn Leu Gln Ala Ile Gln Lys Glu Phe Ser Glu Ile Arg Asp Ser Val Gln Ala Glu Asp Thr Asn Ile Asp Ile Arg Ile 55 Leu Arg Thr Thr Glu Ser Leu Lys Asp Ile Lys Ser Leu Asp Arg Cys Cys Phe Leu Arg His Leu Val Arg Phe Tyr Leu Asp Arg Val Phe Lys Val Tyr Gln Thr Pro Asp His His Thr Leu Arg Lys Ile Ser Ser Leu 100 105 Ala Asn Ser Phe Leu Ile Ile Lys Lys Asp Leu Ser Val Cys His Ser 120 His Met Ala Cys His Cys Gly Glu Glu Ala Met Glu Lys Tyr Asn Gln 130 135 140 Ile Leu Ser His Phe Ile Glu Leu Glu Leu Gln Ala Ala Val Lys 145 150 155 Ala Leu Gly Glu Leu Gly Ile Leu Leu Arg Trp Met Glu Glu Met Leu 165 170

<210> 6

<211> 152

<212> PRT

<213> Mus musculis

<400> 6

Leu Lys Thr Leu His Leu Gly Ser Cys Val Ile Thr Ala Asn Leu Gln 1 5 10 15 15 Ala Ile Gln Lys Glu Phe Ser Glu Ile Arg Asp Ser Val Gln Ala Glu 20 25 30 Asp Thr Asn Ile Asp Ile Arg Ile Leu Arg Thr Thr Glu Ser Leu Lys 35 40 45 Asp Ile Lys Ser Leu Asp Arg Cys Cys Phe Leu Arg His Leu Val Arg 50 55 60 Phe Tyr Leu Asp Arg Val Phe Lys Val Tyr Gln Thr Pro Asp His His 65 70 75 80

```
Thr Leu Arg Lys Ile Ser Ser Leu Ala Asn Ser Phe Leu Ile Ile Lys
                                    90
Lys Asp Leu Ser Val Cys His Ser His Met Ala Cys His Cys Gly Glu
                                105
            100
                                                    110
Glu Ala Met Glu Lys Tyr Asn Gln Ile Leu Ser His Phe Ile Glu Leu
                            120
Glu Leu Gln Ala Ala Val Val Lys Ala Leu Gly Glu Leu Gly Ile Leu
                        135
                                            140
Leu Arg Trp Met Glu Glu Met Leu
                    150
      <210> 7
      <211> 144
      <212> PRT
      <213> Mus musculis
      <400> 7
Cys Val Ile Thr Ala Asn Leu Gln Ala Ile Gln Lys Glu Phe Ser Glu
                                    10
Ile Arg Asp Ser Val Gln Ala Glu Asp Thr Asn Ile Asp Ile Arg Ile
                                25
Leu Arg Thr Thr Glu Ser Leu Lys Asp Ile Lys Ser Leu Asp Arg Cys
Cys Phe Leu Arg His Leu Val Arg Phe Tyr Leu Asp Arg Val Phe Lys
Val Tyr Gln Thr Pro Asp His His Thr Leu Arg Lys Ile Ser Ser Leu
                    70
                                        75
Ala Asn Ser Phe Leu Ile Ile Lys Lys Asp Leu Ser Val Cys His Ser
                                     90
His Met Ala Cys His Cys Gly Glu Glu Ala Met Glu Lys Tyr Asn Gln
            100
                                 105
Ile Leu Ser His Phe Ile Glu Leu Glu Leu Gln Ala Ala Val Val Lys
                             120
                                                 125
        115
Ala Leu Gly Glu Leu Gly Ile Leu Leu Arg Trp Met Glu Glu Met Leu
    130
                         135
                                             140
      <210> 8
      <211> 154
      <212> PRT
```

<213> Mus musculis

<400> 8

Met Lys Gly Phe Gly Leu Ala Phe Gly Leu Phe Ser Ala Val Gly Phe Leu Leu Trp Thr Pro Leu Thr Gly Leu Lys Thr Leu His Leu Gly Ser 25 Cys Val Ile Thr Ala Asn Leu Gln Ala Ile Gln Lys Glu Phe Ser Glu Ile Arg Asp Ser Val Ser Leu Asp Arg Cys Cys Phe Leu Arg His Leu Val Arg Phe Tyr Leu Asp Arg Val Phe Lys Val Tyr Gln Thr Pro Asp His His Thr Leu Arg Lys Ile Ser Ser Leu Ala Asn Ser Phe Leu Ile 90 Ile Lys Lys Asp Leu Ser Val Cys His Ser His Met Ala Cys His Cys 100 105 Gly Glu Glu Ala Met Glu Lys Tyr Asn Gln Ile Leu Ser His Phe Ile 120 125 Glu Leu Glu Leu Gln Ala Ala Val Val Lys Ala Leu Gly Glu Leu Gly 135 140 Ile Leu Leu Arg Trp Met Glu Glu Met Leu 145 150

<210> 9

<211> 130

<212> PRT

<213> Homo sapiens

<400> 9

Leu Lys Thr Leu His Leu Gly Ser Cys Val Ile Thr Ala Asn Leu Gln Ala Ile Gln Lys Glu Phe Ser Glu Ile Arg Asp Ser Val Ser Leu Asp 25 Arg Cys Cys Phe Leu Arg His Leu Val Arg Phe Tyr Leu Asp Arg Val 40 Phe Lys Val Tyr Gln Thr Pro Asp His His Thr Leu Arg Lys Ile Ser 60 Ser Leu Ala Asn Ser Phe Leu Ile Ile Lys Lys Asp Leu Ser Val Cys His Ser His Met Ala Cys His Cys Gly Glu Glu Ala Met Glu Lys Tyr 85 90 Asn Gln Ile Leu Ser His Phe Ile Glu Leu Glu Leu Gln Ala Ala Val 105 Val Lys Ala Leu Gly Glu Leu Gly Ile Leu Leu Arg Trp Met Glu Glu 115 120 125

Met	130									•						
	<2 <2	212>	3516		oiens	5										
	<2	220> 221> 222>	CDS (237	')	(189	95)										
ctgg gcgt	agcto gggaa cggca	agc o	taged etggg eagag	ggad gcccd	eg co	gcto gcgco	gtggo ggago	gga tga	aggco aggco	gccc cgcg	tggg cggd	gacto ccgco	cag g	gtcgo tggco	tggttt cctgga cccagc cc atg Met 1	60 120 180 239
			ggc Gly 5				_		_	-	_	_	_	_	-	287
			ctc Leu													335
			ttg Leu				-									383
_	_		gtc Val												_	431
	_		tac Tyr		-	_							_			479
			tca Ser 85			-								-		527

	gct Ala											-		_	_	į	575
	att Ile 115															(523
	cct Pro															(671
	gat Asp															;	719
	aga Arg															ì	767
	ctg Leu															{	815
	tcc Ser 195														gag Glu		863
	aac Asn				Cýs	Val		Val								Ć	911
_	cgc Arg													_		Ċ	959
	caa Gln															10	007
ccc	ata	tct	att	acc	gtg	ttt	ctt	ttt	tct	gtg	atg	ggc	tat	tcc	atc	10)55

WO 01/46261 9

Pro	Ile	Ser 260	Ile	Thr	Val	Phe	Leu 265	Phe	Ser	Val	Met	G1y 270	Tyr	Ser	Ile	
	_	tat Tyr				-										1103
_		tat Tyr			-		-		_					_		1151
		gtg Val									_	-	_			1199
		cat His	_	-	_	_		_			-	-	-	-		1247
_		aat Asn 340	_				-									1295
-		gag Glu										Leu				1343
	_	gac Asp		•	_		_	_						_		1391
-		ctc Leu	-	_				_	-	Lys		_		-		1439
-		gat Asp	-	Arg					Cys					Glu		1487
		agt Ser 420						Ser					Leu			1535

tcg cag gca gcg ttg gca gtc ttg ggc ccg caa acg tta cag tac tca Ser Gln Ala Ala Leu Ala Val Leu Gly Pro Gln Thr Leu Gln Tyr Ser 435 440 445	583
tac acc cct cag ctc caa gac tta gac ccc ctg gcg cag gag cac aca Tyr Thr Pro Gln Leu Gln Asp Leu Asp Pro Leu Ala Gln Glu His Thr 450 465 460 465	631
gac tcg gag gag ggg ccg gag gaa gag cca tcg acg acc ctg gtc gac Asp Ser Glu Glu Gly Pro Glu Glu Glu Pro Ser Thr Thr Leu Val Asp 470 475 480	679
tgg gat ccc caa act ggc agg ctg tgt att cct tcg ctg tcc agc ttc Trp Asp Pro Gln Thr Gly Arg Leu Cys Ile Pro Ser Leu Ser Ser Phe 485 490 495	727
gac cag gat tca gag ggc tgc gag cct tct gag ggg gat ggg ctc gga Asp Gln Asp Ser Glu Gly Cys Glu Pro Ser Glu Gly Asp Gly Leu Gly 500 505 510	775
gag gag ggt ctt cta tct aga ctc tat gag gag ccg gct cca gac agg Glu Glu Gly Leu Leu Ser Arg Leu Tyr Glu Glu Pro Ala Pro Asp Arg 515 520 525	823
cca cca gga gaa aat gaa acc tat ctc atg caa ttc atg gag gaa tgg Pro Pro Gly Glu Asn Glu Thr Tyr Leu Met Gln Phe Met Glu Glu Trp 530 535 540 545	871
ggg tta tat gtg cag atg gaa aac tgatgccaac acttcctttt gccttttgtt 19 Gly Leu Tyr Val Gln Met Glu Asn 550	925
	985
	045 105
gtccaggggc tggccggttg ttctatgcag agaaagcagt caataaatgt ttgccagact 21	165
The state of the s	225
	285 345
	405
	465
	525 585

WO 01/46261

11

tttagtaggg gcttattgtc tctcaaaact aacctaaaag aaaatgacac attttataat 2645 agaatattac atttatttct ggaagtgtgt tttcaaaaag atatttacat agtctgtaaa 2705 ctagaaagtg ttaggtaaag ctctaggtta ctgtgttact attataatat taaacattcg 2765 aataggcagt cgttcaaaga ctctttggaa tatctatgaa tgaatatcct ctattcttat 2825 aatattaaaa cccataagta aatataggac atacaagaga aatgagttaa atgactatgt 2885 2945 caatagcttt ctagttcatt tccagtaact gttcccatct cctttaccac ttgttaagaa 3005 aattaaattc ttcagtcacg ctgctttaaa atgggacaaa atctattaag ttgaaccata 3065 tataattgtg gatatttggc tgtttttaat ctgacaagca gtaacttcat atggtttgcc 3125 ttaatatata tttgttttag tcatgaactc ataatccatt gatgctcttt catgagaaga 3185 gatatgaccc atatttcctt attgatatta ttggtacagg cagacaaccc tggtaggaga 3245 gatggattct ggggtcatga cctttcgtga ttatccgcaa atgcaaacag tttcagatct 3305 aatggtttaa tttagggagt aattatatta atcagagtgt tctgttattc tcaatcttta 3365 tagaaacgat tetgetggtt ttgaagaaca gatgtattac actaactgta aaagtagtte 3425 aagagtgaga aagaataaat tgttattaag agcaaaagaa aaataaagtg attgatgata 3485 aaaaaaaaa aaaaaaagcg gccgcctcga g 3516

<210> 11

<211> 553

<212> PRT

<213> Homo sapiens

<400> 11

Met Arg Ala Pro Gly Arg Pro Ala Leu Arg Pro Leu Pro Leu Pro Pro 1 5 10 Leu Leu Leu Leu Leu Ala Ala Pro Trp Gly Arg Ala Val Pro Cys 25 Val Ser Gly Gly Leu Pro Lys Pro Ala Asn Ile Thr Phe Leu Ser Ile 35 40 45 Asn Met Lys Asn Val Leu Gln Trp Thr Pro Pro Glu Gly Leu Gln Gly 55 60 Val Lys Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly Gln Lys Lys 70 75 Trp Leu Asn Lys Ser Glu Cys Arg Asn Ile Asn Arg Thr Tyr Cys Asp 85 90 Leu Ser Ala Glu Thr Ser Asp Tyr Glu His Gln Tyr Tyr Ala Lys Val 100 105 Lys Ala Ile Trp Gly Thr Lys Cys Ser Lys Trp Ala Glu Ser Gly Arg 125 115 120 Phe Tyr Pro Phe Leu Glu Thr Gln Ile Gly Pro Pro Glu Val Ala Leu 130 135 140 Thr Thr Asp Glu Lys Ser Ile Ser Val Val Leu Thr Ala Pro Glu Lys 145 150 155 160

WO 01/46261 PCT/US00/35305 12

				165)				170)				175	Tyr
			T8()				185	,				190)	i Arg
		195)				200					205)		Leu
	210)				215					220	1			Gly
225	•				230					235	,				Leu 240
				245					250	1				255	Va1
			260					265					270	1	Ser
		275					280					285			Leu
	290				Asn	295					300				
305					Asn 310					315					320
				325	Asp				330					335	
			340		Pro			345					350		
		355			Lys		360					365			
	3/0				G1u	375					380				
385					Arg 390					395					400
				405	Arg				410					415	
			420		Gln			425					430		
		435			Leu		440					445			
	450					455					460				
465					Gly 470					475					4 80
Asp	Irp	Asp	Pro	G1n 485	Thr	Gly ,	Arg		Cys 490	Ile	Pro	Ser	Leu	Ser 495	Ser

```
Phe Asp Gln Asp Ser Glu Gly Cys Glu Pro Ser Glu Gly Asp Gly Leu
                                505
            500
Gly Glu Glu Gly Leu Leu Ser Arg Leu Tyr Glu Glu Pro Ala Pro Asp
                            520
Arg Pro Pro Gly Glu Asn Glu Thr Tyr Leu Met Gln Phe Met Glu Glu
                        535
                                            540
Trp Gly Leu Tyr Val Gln Met Glu Asn
545
                    550
      <210> 12
      <211> 221
      <212> PRT
      <213> Homo sapiens
      <400> 12
Val Pro Cys Val Ser Gly Gly Leu Pro Lys Pro Ala Asn Ile Thr Phe
Leu Ser Ile Asn Met Lys Asn Val Leu Gln Trp Thr Pro Pro Glu Gly
                                25
Leu Gln Gly Val Lys Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly
                            40
Gln Lys Lys Trp Leu Asn Lys Ser Glu Cys Arg Asn Ile Asn Arg Thr
Tyr Cys Asp Leu Ser Ala Glu Thr Ser Asp Tyr Glu His Gln Tyr Tyr
                    70
                                        75
Ala Lys Val Lys Ala Ile Trp Gly Thr Lys Cys Ser Lys Trp Ala Glu
                                    90
Ser Gly Arg Phe Tyr Pro Phe Leu Glu Thr Gln Ile Gly Pro Pro Glu
            100
                                105
Val Ala Leu Thr Thr Asp Glu Lys Ser Ile Ser Val Val Leu Thr Ala
                            120
Pro Glu Lys Trp Lys Arg Asn Pro Glu Asp Leu Pro Val Ser Met Gln
    130
                        135
                                             140
Gln Ile Tyr Ser Asn Leu Lys Tyr Asn Val Ser Val Leu Asn Thr Lys
                    150
                                         155
Ser Asn Arg Thr Trp Ser Gln Cys Val Thr Asn His Thr Leu Val Leu
                165
                                    170
Thr Trp Leu Glu Pro Asn Thr Leu Tyr Cys Val His Val Glu Ser Phe
            180
                                185
                                                     190
Val Pro Gly Pro Pro Arg Arg Ala Gln Pro Ser Glu Lys Gln Cys Ala
                            200
Arg Thr Leu Lys Asp Gln Ser Ser Glu Phe Lys Ala Lys
    210
                        215
                                             220
```

WO 01/46261 PCT/US00/35305

<210> 13 <211> 971 <212> DNA <213> Homo sapiens														
<220> <221> CDS <222> (18)(950)														
<pre><400> 13 gaattcgagt ctaccaa atg cag act ttc aca atg gtt cta gaa gaa atc</pre>														
tgg aca agt ctt ttc atg tgg ttt ttc tac gca ttg att cca tgt ttg Trp Thr Ser Leu Phe Met Trp Phe Phe Tyr Ala Leu Ile Pro Cys Leu 15 20 25	98													
ctc aca gat gaa gtg gcc att ctg cct gcc cct cag aac ctc tct gta Leu Thr Asp Glu Val Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val 30 35 40	146													
ctc tca acc aac atg aag cat ctc ttg atg tgg agc cca gtg atc gcg Leu Ser Thr Asn Met Lys His Leu Leu Met Trp Ser Pro Val Ile Ala 45 50 55	194													
cct gga gaa aca gtg tac tat tct gtc gaa tac cag ggg gag tac gag Pro Gly Glu Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu 60 65 70 75	242													
agc ctg tac acg agc cac atc tgg atc ccc agc agc tgg tgc tca ctc Ser Leu Tyr Thr Ser His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu 80 85 90	290													
act gaa ggt cct gag tgt gat gtc act gat gac atc acg gcc act gtg Thr Glu Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val 95 100 105	338													
cca tac aac ctt cgt gtc agg gcc aca ttg ggc tca cag acc tca gcc Pro Tyr Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala 110 115 120	386													

WO 01/46261 PCT/US00/35305

	-		ctg Leu													434
_			atg Met		•			-				_	-			482
			ctg Leu													530
			ggt Gly 175							_			_			578
			cac His								_	-		-		626
			aca Thr									. –	_		_	674
			tgt Cys							_			_	-	_	722
			gcc Ala	Phe		Gly	Phe	Met	Leu	Ile						770
_		-	tgg Trp 255		_			_		_			_	_		818
			ctc Leu		_		_							_	_	866
			tgc Cys													914

-	tct Ser		-	-								taggtttgcg				9	960
gaag	gcto	ga g]													9	971
	<2 <2	210> 211> 212> 213>	311	sap	oiens	3											
	</td <td>100></td> <td>14</td> <td></td>	100>	14														
Met 1	Gln	Thr	Phe	Thr 5	Met	Val	Leu	Glu	Glu 10	Ile	Trp	Thr	Ser	Leu 15	Phe		
	Trp	Phe	Phe 20		Ala	Leu	Ile	Pro 25		Leu	Leu	Thr	Asp 30		Val		
Ala	Ile	Leu 35	Pro	Ala	Pro	Gln	Asn 40	Leu	Ser	Val	Leu	Ser 45	Thr	Asn	Met		
Lys	His 50		Leu	Met	Trp	Ser 55	. –	Val	Ile	Ala	Pro 60		Glu	Thr	Val		
Tyr 65	Tyr	Ser	Val	Glu	Tyr 70		Gly	Glu	Tyr	G1u 75	Ser	Leu	Tyr	Thr	Ser 80		
	Ile	Trp	Ile	Pro 85		Ser	Trp	Cys	Ser 90		Thr	Glu	Gly	Pro 95	Glu		
Cys	Asp	Val	Thr 100		Asp	Ile	Thr	Ala 105	Thr	Val	Pro	Tyr	Asn 110	Leu	Arg		
Val	Arg	Ala 115	Thr	Leu	Gly	Ser	Gln 120	Thr	Ser	Ala	Trp	Ser 125	Пe	Leu	Lys		
His	Pro 130		Asn	Arg	Asn	Ser 135		Ile	Leu	Thr	Arg 140	Pro	Gly	Met	Glu		
Ile 145	Thr	Lys	Asp	Gly	Phe 150		Leu	Val	Пe	G1u 155	Leu	Glu	Asp	Leu	Gly 160		
	Gln	Phe	Glu	Phe 165		Val	Ala	Tyr	Trp 170			Glu	Pro	Gly 175			
Glu	Glu	His	Val 180		Met	Val	Arg	Ser 185	Gly	Gly	Ile	Pro	Val 190		Leu		
Glu	Thr	Met 195	Glu	Pro	Gly	Ala	A1a 200	Tyr		Val	Lys	Ala 205	Gln	Thr	Phe		
Val	Lys 210			Gly	Arg	Tyr 215	Ser		Phe	Ser	G1n 220		Glu	Cys	Val		
G1u 225	Val	Gln	Gly	Glu	A1a 230			Leu	Val	Leu 235	Ala	Leu	Phe	Ala	Phe 240		

Val Gly Phe Met Leu Ile Leu Val Val Val Pro Leu Phe Val Trp Lys 250 Met Gly Arg Leu Leu Gln Tyr Ser Cys Cys Pro Val Val Leu Pro 260 265 270 Asp Thr Leu Lys Ile Thr Asn Ser Pro Gln Lys Leu Ile Ser Cys Arg 280 Arg Glu Glu Val Asp Ala Cys Ala Thr Ala Val Met Ser Pro Glu Glu 295 300 Leu Leu Arg Ala Trp Ile Ser 305 310 <210> 15 <211> 203 <212> PRT <213> Homo sapiens <400> 15 Asp Glu Val Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser 10 Thr Asn Met Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly 25 Glu Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu 40 Tyr Thr Ser His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr 70 75 Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser 90 Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro 105 Gly Met Glu Ile Thr Lys Asp Gly Phe His Leu Val Ile Glu Leu Glu 120 125 115 Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Arg Arg Glu 135 140 Pro Gly Ala Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro 150 155 Val His Leu Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala 165 170 Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr 180 190 185 Glu Cys Val Glu Val Gln Gly Glu Ala Ile Pro 195 200

<210> 16 <211> 33 <212> DNA <213> Homo sapiens	
<400> 16 gcgaattcga gtctaccaaa tgcagacttt cac	33
<210> 17 <211> 32 <212> DNA ' <213> Homo sapiens	
<400> 17 cgctcgagcc ttccgcaaac ctatgagatc ca	32
<210> 18 <211> 1379 <212> DNA <213> Homo sapiens	
<220> <221> CDS <222> (132)(1034)	
<pre><400> 18 tcgacccacg cgtccgcgct gcgactcaga cctcagctcc aacatatgca ttctgaagaa agatggctga gatggacaga atgctttatt ttggaaagaa acaatgttct aggtcaaact gagtctacca a atg cag act ttc aca atg gtt cta gaa gaa atc tgg aca</pre>	60 120 170
Met Gln Thr Phe Thr Met Val Leu Glu Glu Ile Trp Thr 1 5 10	
agt ctt ttc atg tgg ttt ttc tac gca ttg att cca tgt ttg ctc aca Ser Leu Phe Met Trp Phe Phe Tyr Ala Leu Ile Pro Cys Leu Leu Thr 15 20 25	218
gat gaa gtg gcc att ctg cct gcc cct cag aac ctc tct gta ctc tca Asp Glu Val Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser 30 35 40 45	266
acc aac atg aag cat ctc ttg atg tgg agc cca gtg atc gcg cct gga	314

Thr	Asn	Met	Lys	His 50	Leu	Leu	Met	Trp	Ser 55	Pro	Val	Ile	Ala	Pro 60	Gly	,	
		gtg Val															362
	_	agc Ser 80						-	_		-				_		410
		gag Glu	_	-	_		_	-			-						458
		cgt Arg															506
	_	aag Lys					_							_			554
	-	gag Glu								-	-			_			602
-	_	ggg Gly 160		_						-			_				650
		gcc Ala		-		-		_			_						698
		cta Leu															746
		ttc Phe	_	_	Ala					Ser							794

gaa tgt gtg gag gtg caa gga gag gcc att ccc ctg gta Glu Cys Val Glu Val Gln Gly Glu Ala Ile Pro Leu Val 225 230	
ttt gcc ttt gtt ggc ttc atg ctg atc ctt gtg gtc gtg Phe Ala Phe Val Gly Phe Met Leu Ile Leu Val Val Val 240 245 250	
gtc tgg aaa atg ggc cgg ctg ctc cag tac tcc tgt tgc Val Trp Lys Met Gly Arg Leu Leu Gln Tyr Ser Cys Cys 255 260 265	
gtc ctc cca gac acc ttg aaa ata acc aat tca ccc cag Val Leu Pro Asp Thr Leu Lys Ile Thr Asn Ser Pro Gln 270 275 280	•
ctg cag aag gga gga ggt gga tgc ctg tgc cac ggc tgt Leu Gln Lys Gly Gly Gly Gly Cys Leu Cys His Gly Cys 290 295	_
tgaggaactc ctcagggcct ggatctcata tcaggtttgc ggaagggcggagaacctgg tctgcatgac atggaaacca tgaggggaca agttgtgt ccacggacaa gggatgagag aagtaggaag agcctgttgt ctacaagtactagaggca gggtggtttg tctaacagaa caactgactg aggctatgtctagacttt gggcttccac ttgcttggct gagcaaccct gggaaaaccttcggtccca agttttctca tctgtaatgg gggatcccta caaaactg	ttt ctgttttccg 1154 tct agaagcaacc 1214 ggg ggttgtgacc 1274 gtg acttcatccc 1334
<210> 19 <211> 301 <212> PRT <213> Homo sapiens	
<pre><400> 19 Met Gln Thr Phe Thr Met Val Leu Glu Glu Ile Trp Thr</pre>	Ser Leu Phe
1 5 10 Met Trp Phe Phe Tyr Ala Leu Ile Pro Cys Leu Leu Thr	15 Asp Glu Val
20 25 Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser	30 Thr Asn Met
35 40 45 Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly	Glu Thr Val
50 55 60	
Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu	lyr inr Ser

```
His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu Gly Pro Glu
Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr Asn Leu Arg
                                105
Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser Ile Leu Lys
                            120
His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro Gly Met Glu
                        135
                                            140
Ile Pro Lys His Gly Phe His Leu Val Ile Glu Leu Glu Asp Leu Gly
                    150
                                        155
Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Thr Arg Glu Pro Gly Ala
                                    170
                165
Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro Val His Leu
                                185
            180
                                                    190
Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala Gln Thr Phe
                            200
Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr Glu Cys Val
                        215
Glu Val Gln Gly Glu Ala Ile Pro Leu Val Leu Ala Leu Phe Ala Phe
225
                    230
                                        235
                                                             240
Val Gly Phe Met Leu Ile Leu Val Val Val Pro Leu Phe Val Trp Lys
                245
                                    250
Met Gly Arg Leu Leu Gln Tyr Ser Cys Cys Pro Val Val Leu Pro
            260
                                265
Asp Thr Leu Lys Ile Thr Asn Ser Pro Gln Val Asn Gln Leu Gln Lys
        275
                            280
                                                285
Gly Gly Gly Cys Leu Cys His Gly Cys Asp Val Ser
    290
                        295
                                            300
      <210> 20
      <211> 1081
      <212> DNA
      <213> Homo sapiens
      <220>
      <221> CDS
      <222> (9)...(1067)
      <400> 20
ggccggcc atg cag act ttc aca atg gtt cta gaa gaa atc tgg aca agt
```

Met Gln Thr Phe Thr Met Val Leu Glu Glu Ile Trp Thr Ser

							gca Ala				_	-			_	98	
							cct Pro					_				146	
							tgg Trp									194	
							tac Tyr 70						_	_		242	
	-						agc Ser	_		-				_		290	
							gac Asp		_	_						338	
	-	_				_	ggc Gly		_			_		_		386	
			Pro	Phe	Asn	Arg	aac Asn	Ser	Thr	Пe		Thr	Arg	Pro		434	
_							ttc Phe 150		_	-			_		_	482	
							ctt Leu		-			-				530	
							atg Met									578	

PCT/US00/35305

				atg Met 195									_	_	_	626
		-	-	gcc Ala					-	_		_	-		_	674
				caa Gln								_				722
				ggt Gly		-	-			-	-			-		770
		-		tct Ser	_		-	-					-		-	818
				aat Asn 275					-		_		-	_		866
		_		gcc Ala			_				-		_	_		914
	-	-		aag Lys	-	_			_		-	_		_	_	962
_	_		_	gac Asp						_		_	_	-	-	1010
		_		ctg Leu	_	_		-		_	_				gga Gly 350	1058
	tgt Cys		tct	agag	gcg (cgcc										1081

<210> 21 <211> 352 <212> PRT <213> Homo sapiens

<400> 21 Met Gln Thr Phe Thr Met Val Leu Glu Glu Ile Trp Thr Ser Leu Phe Met Trp Phe Phe Tyr Ala Leu Ile Pro Cys Leu Leu Thr Asp Glu Val 25 Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser Thr Asn Met Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly Glu Thr Val 60 Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu Tyr Thr Ser 75 70 His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu Gly Pro Glu 90 Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr Asn Leu Arg 105 Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser Ile Leu Lys 120 His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro Gly Met Glu 135 Ile Pro Lys His Gly Phe His Leu Val Ile Glu Leu Glu Asp Leu Gly 155 150 Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Thr Arg Glu Pro Gly Ala 175 170 165 Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro Val His Leu 185 180 Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala Gln Thr Phe 205 200 Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr Glu Cys Val 220 215 Glu Val Gln Gly Glu Ala Gly Gly Gly Gly Ser Gly Gly Gly Ser 235 230 Gly Gly Gly Ser Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe 250 245 Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys 265 Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val 280 275

Asp Asn A	la Leu	Gln	Ser	G1y 295	Asn	Ser	G1n	Glu	Ser 300	Val	Thr	Glu	Gln	
Asp Ser L	ys Asp	Ser	Thr 310		Ser	Leu	Ser	Ser 315		Leu	Thr	Leu	Ser 320	
Lys Ala A	sp Tyr	G1u 325	Lys	His	Lys	Val	Tyr 330	Ala	Cys	Glu	Val	Thr 335	His	
Gln Gly L	eu Ser 340		Pro	Val	Thr	Lys 345		Phe	Asn	Arg	Gly 350		Cys	
<21 <21	0> 22 1> 180 2> DNA 3> Hom		oiens	5										
	0> 1> CDS 2> (8)		1789))										
gtcgacc a M	0> 22 tg gat et Asp 1	_	_	_	_			_	_		_	_	_	49
tgt ggc g Cys Gly A 15														.97
aga cgc t Arg Arg P	_	-	-		-	-				_				145
gca aac a Ala Asn I		Phe					_	-		-				193
act cca c Thr Pro P						-							-	241
tat ttc a Tyr Phe I 80										Ser				289

			aga Arg			_	-			-	_			_		337
			tat Tyr											-		385
			gct Ala 130	_	-								-			433
			cca Pro			-	_			_		_				481
_	-	-	aca Thr	-			_		_	_			•	-		529
			atg Met													577
			act Thr													625
			gtg Val 210	Leu	Thr	Trp	_	Glu	Pro			Leu		Cys	-	673
			tcc Ser		-					_	_	-	_			721
	_	_	tgt Cys	_			_		-							769
			ggc Gly											_		817

WO 01/46261 PCT/US00/35305

tcg (Ser	-			-	-					-						865
gcg (_	_		-	_	-	-	-				-	-		_	913
gtg Val	_					_	_		-						_	, 961
gct Ala	_		_								-	_				1009
gtg Val 335			_	_	-			_				_				1057
cac His	_		_			-		_	_		_					1105
tgt Cys	-					-							-			1153
ggg Gly	-	_	Ser	_			Phe		Pro	Lys	Pro	_	Asp			1201
atg Met							-		_				_		_	1249
cac His 415	_	_			_	_						-				1297
gtg	cat	aat	gcc	aag	aca	aag	ccg	cgg	gag	gag	cag	tac	aac	agc	acg	1345

WO 01/46261 PCT/US00/35305

Val	His	Asn	Ala	Lys 435	Thr	Lys	Pro	Arg	G1u 440	G1u	Gln	Tyr	Asn	Ser 445	Thr	
					_			_	_		_	_		ctg Leu		1393
											_			tcc Ser		1441
														cca Pro		1489
														cag Gln		1537
														gcc Ala 525		1585
							_					_		acg Thr		1633
												-	_	ctc Leu		1681
					-	-	-			-			-	tcc Ser	-	1729
														tcc Ser		1777
	ccg Pro			taat	ctag	gat c	:t									1801

<210> 23

<211> 594 <212> PRT <213> Homo sapiens

<400> 23 Met Asp Ala Met Lys Arg Gly Leu Cys Cys Val Leu Leu Cys Gly Ala Val Phe Val Ser Leu Ser Gln Glu Ile His Ala Glu Leu Arg Arg 25 Phe Arg Arg Val Pro Cys Val Ser Gly Gly Leu Pro Lys Pro Ala Asn 40 Ile Thr Phe Leu Ser Ile Asn Met Lys Asn Val Leu Gln Trp Thr Pro 55 Pro Glu Gly Leu Gln Gly Val Lys Val Thr Tyr Thr Val Gln Tyr Phe 75 Ile Tyr Gly Gln Lys Lys Trp Leu Asn Lys Ser Glu Cys Arg Asn Ile 90 85 Asn Arg Thr Tyr Cys Asp Leu Ser Ala Glu Thr Ser Asp Tyr Glu His 110 105 Gin Tyr Tyr Ala Lys Val Lys Ala Ile Trp Gly Thr Lys Cys Ser Lys 120 Trp Ala Glu Ser Gly Arg Phe Tyr Pro Phe Leu Glu Thr Gln Ile Gly 140 135 Pro Pro Glu Val Ala Leu Thr Thr Asp Glu Lys Ser Ile Ser Val Val 155 150 Leu Thr Ala Pro Glu Lys Trp Lys Arg Asn Pro Glu Asp Leu Pro Val 170 Ser Met Gln Gln Ile Tyr Ser Asn Leu Lys Tyr Asn Val Ser Val Leu 185 Asn Thr Lys Ser Asn Arg Thr Trp Ser Gln Cys Val Thr Asn His Thr 200 Leu Val Leu Thr Trp Leu Glu Pro Asn Thr Leu Tyr Cys Val His Val 215 Glu Ser Phe Val Pro Gly Pro Pro Arg Arg Ala Gln Pro Ser Glu Lys 235 230 Gln Cys Ala Arg Thr Leu Lys Asp Gln Gly Gly Gly Ser Gly Gly 250 Gly Gly Ser Gly Gly Gly Ser Ala Ser Thr Lys Gly Pro Ser Val 265 260 Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala 285 280 Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser 300 295 290

PCT/US00/35305 WO 01/46261 **3**0

305					310					His 315					320
				325					330	Ser				335	
			340					345		Cys			350		
Pro	Ser	Asn 355	Thr	Lys	Val	Asp	Lys 360	Lys	Val	Glu	Pro	Lys 365	Ser	Cys	Asp
Lys	Thr 370	His	Thr	Cys	Pro	Pro 375	Cys	Pro	Ala	Pro	G1u 380	Ala	Glu	Gly	Ala
Pro 385		Val	Phe	Leu	Phe 390	Pro	Pro	Lys	Pro	Lys 395	Asp	Thr	Leu	Met	I1e 400
Ser	Arg	Thr	Pro	G1u 405		Thr	Cys	Val	Val 410	Val	Asp	Val	Ser	His 415	Glu
Asp	Pro	Glu	Val 420		Phe	Asn	Trp	Tyr 425		Asp	G1y	Val	G1u 430	Val	His
Asn	Ala	Lys 435		Lys	Pro	Arg	G1u 440		Gln	Tyr	Asn	Ser 445		Tyr	Arg
	450					455				Asp	460				
G1u 465		· Lys	Cys	Lys	Val 470		Asn	Lys	Ala	Leu 475		Ser	Ser	Ile	G1u 480
Lys	Thr	· Ile	Ser	Lys 485		Lys	Gly	Gln	Pro 490	Arg	Glu	Pro	Gln	Va1 495	Tyr
Thr	Leu	ı Pro	Pro 500		· Arg	Asp	Glu	Leu 505		Lys	Asn	Gln	Val 510		Leu
Thr	Cys	Leu 515		Lys	Gly	Phe	Tyr 520		Ser	· Asp	lle	A1 a 525		Glu	Trp
	530)				535)				540)			Val
Leu 545		Ser	· Ast	Gly	/ Ser 550		Phe	e Leu	ı Tyr	Ser 555		: Lei	ı Thr	· Val	Asp 560
				565	5				57()				575	
G1ı	u Ala	a Lei	u His 580		n His	Tyr	⁻ Thr	Glr 585		s Ser	^ Leu	ı Ser	• Let 590	ı Ser)	· Pro
Gly	y Lys	S													

<210> 24

<211> 29

<212> DNA

<213> Homo sapiens

<400> 24	
ggccggccat gcagactttc acaatggtt	29
<210> 25	
<211> 52	
<212> DNA	
<213> homo sapiens	
<400> 25	
teegetaceg eegecteeac tgecaceace teeggeetet eettgeacet ee	52
<210> 26	
<211> 53	
<212> DNA	
<213> Homo sapiens	
<400> 26	
gtggaggcgg cggtagcgga ggcggtggca gtcgaactgt ggctgcacca tct	53
<210> 27	
<211> 38	
<212> DNA	
<213> Homo sapiens	
<400> 27	
ggcgcgcctc tagattaaca ctctcccctg ttgaagct	38
<210> 28	
<211> 30	
<212> DNA	
<213> Homo sapiens	
<400> 28	
gtcgaccatg gatgcaatga agagagggct	30
<210> 29	
<211> 30	
<212> DNA	
<213> Homo sapiens	
<400> 29	
cacagggaac tctacggaag cgtctcaact	30

.

<210> 30 <211> 33 <212> DNA <213> Homo sapiens	
<400> 30 cttccgtaga gttccctgtg tctctggtgg ttt	33
<210> 31 <211> 53 <212> DNA <213> Homo sapiens	
<pre><400> 31 gccagagcca cctccgcctg aaccgcctcc accttgatct ttcaaagtcc tgg</pre>	53
<210> 32 <211> 51 <212> DNA <213> Homo sapiens	
<400> 32 caggcggagg tggctctggc ggtggcggat cggcctccac caagggccca t	51
<210> 33 <211> 20 <212> DNA <213> Homo sapiens	
<400> 33	20
<pre>ctgggcacgg tgggcatgtg</pre>	20
<400> 34 cacatgccca ccgtgcccag	20
<210> 35 <211> 31 <212> DNA	

<213> Homo sapiens <400> 35 31 agatctagat tatttacccg gagacaggga g <210> 36 <211> 1806 <212> DNA <213> Mus musculus <220> <221> CDS <222> (38)...(1675) <400> 36 cgccgcgttc ccgagatgtg acccgaactg acagccc atg cac act ccc ggg acc 55 Met His Thr Pro Gly Thr 1 5 ccg gcg ccg ggc cac ccg gac ccg ccg cca ctg ttg ctg ctc acg ctg 103 Pro Ala Pro Gly His Pro Asp Pro Pro Pro Leu Leu Leu Leu Thr Leu 10 15 ctt ctg ctg ctg gcc gct tcg gga cgc gca gtt cct tgt gtc ttc tgt 151 Leu Leu Leu Leu Ala Ala Ser Gly Arg Ala Val Pro Cys Val Phe Cys 25 30 ggt ttg cct aaa cct aca aat atc acc ttc tta tcc atc aac atg aag 199 Gly Leu Pro Lys Pro Thr Asn Ile Thr Phe Leu Ser Ile Asn Met Lys 40 45 50 aat gtc ctg cat tgg aat cca cca gag agt cta cac gga gtt gaa gtc 247 Asn Val Leu His Trp Asn Pro Pro Glu Ser Leu His Gly Val Glu Val 55 60 65 70 aca tac act gtg caa tat ttc ata tat ggg cag aag aaa tgg ctg aat 295 Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly Gln Lys Lys Trp Leu Asn 75 80 gcc tct aaa tgc ggg agt atc aac agg acc tac tgt gac ctt tct gtt 343

Ala Ser Lys Cys Gly Ser Ile Asn Arg Thr Tyr Cys Asp Leu Ser Val

95

100

gag acc toa gac tat gaa cac cag tto tat gcc aaa gtg aag gcc att Glu Thr Ser Asp Tyr Glu His Gln Phe Tyr Ala Lys Val Lys Ala Ile tgg gaa gcc agg tgc tcc gaa tgg gcc gag acg gaa cgc ttc tat cct Trp Glu Ala Arg Cys Ser Glu Trp Ala Glu Thr Glu Arg Phe Tyr Pro ttc ttg gaa act caa gtc agc cca cca gag att gcc ctg aca act ggc Phe Leu Glu Thr Gln Val Ser Pro Pro Glu Ile Ala Leu Thr Thr Gly gag aag too ato tot att goo otg aca goa coa gag aag tgg aaa aga Glu Lys Ser Ile Ser Ile Ala Leu Thr Ala Pro Glu Lys Trp Lys Arg aat cca caa gac cac act gtt tct atg caa cag ata tac ccc aat ttg Asn Pro Gln Asp His Thr Val Ser Met Gln Gln Ile Tyr Pro Asn Leu aag tac aat gtg tct gtg tat aac act aag tcg aga aga acg tgg tcc Lys Tyr Asn Val Ser Val Tyr Asn Thr Lys Ser Arg Arg Thr Trp Ser cag tgt gtc acc aac agc aca ctg gtc ctc agc tgg ctg gag ccc aac Gln Cys Val Thr Asn Ser Thr Leu Val Leu Ser Trp Leu Glu Pro Asn act ctg tat tgt gtc cac gtg gag tcc ctt gtc cca ggg ccc cct cgc Thr Leu Tyr Cys Val His Val Glu Ser Leu Val Pro Gly Pro Pro Arg ctc ccg atg cct tct cag aag cag tgc atc agt act ttg gaa gtt caa Leu Pro Met Pro Ser Gln Lys Gln Cys Ile Ser Thr Leu Glu Val Gln aca tca gca tgg aag gct aaa gtc atc ttc tgg tat gtc ttc ctc aca Thr Ser Ala Trp Lys Ala Lys Val Ile Phe Trp Tyr Val Phe Leu Thr tct gtt atc gtg ttt ctt ttc tcc gca att ggc tac ttg gtt tac cgt Ser Val Ile Val Phe Leu Phe Ser Ala Ile Gly Tyr Leu Val Tyr Arg

			-	ggc Gly	-	_										919
	_		-	att Ile			-	-		_			_			967
				atc Ile 315			_									1015
		-	-	aat Asn												1063
	-			cac His			_									1111
				tta Leu		-		_							-	1159
	_			aga Arg	-						Thr				tgg Trp 390	1207
		_		atc Ile 395	Pro					Asp					tac Tyr	1255
	_		_	Asp					Gly					Ser	tgt Cys	1303
			Glu	gcg Ala				Glu					Pro		gtg Val	1351
act	tca	gca	aac	ttg	gac	сса	cag	ctt	gaa	gac	cta	cat	cac	ctg	ggt	1399

Thr	Ser 440	Ala	Asn	Leu	Asp	Pro 445	Gln	Leu	Glu	Asp	Leu 450	His	His	Leu	Gly	
														tct Ser		1447
														cct Pro 485		1495
														gag Glu	-	1543
														aac Asn		1591
														cgg Arg		1639
				ggg Gly								tagi	igcc	agg		1685
														-	caggtt ctcatc	1745 1805 1806
	<2 <2	210> 211> 212> 213>	546 PRT	muso	culus	5										
Met		100> Thr		Glv	Thr	Pro	Δla	Pro	Glv	Hic	Pro	Δsn	Pro	Pro	Pro	
1				5					10			•		15		
•			20					25					30	Arg		
Val	Pro	Cys 35	Val	Phe	Cys	G1y	Leu 40	Pro	Lys	Pro	Thr	Asn 45	Ile	Thr	Phe	

Leu	Ser 50	Ile	Asn	Met	Lys	Asn 55	Val	Leu	His	Trp	Asn 60	Pro	Pro	Glu	Ser
Leu 65	His	Gly	Val	Glu	Va1 70	Thr	Tyr	Thr	Va1	G1n 75	Tyr	Phe	Ile	Tyr	Gly 80
Gln	Lys	Lys	Trp	Leu 85	Asn	Ala	Ser	Lys	Cys 90	Gly	Ser	Ile	Asn	Arg 95	Thr
Tyr	Cys	Asp	Leu 100	Ser	Val	Glu	Thr	Ser 105	Asp	Tyr	Glu	His	Gln 110	Phe	Tyr
Ala	Lys	Val 115	Lys	Ala	Ile	Trp	G1u 120	Ala	Arg	Cys	Ser	G1u 125	Trp	Ala	Glu
	130			-		135					140			Pro	
145					150					155				Thr	160
				165					170					Met 175	
Gln	Ile	Tyr	Pro 180	Asn	Leu	Lys	Tyr	Asn 185	Val	Ser	Val	Tyr	Asn 190	Thr	Lys
Ser	Arg	Arg 195	Thr	Trp	Ser	Gln	_	Val		Asn	Ser	Thr 205	Leu	Val	Leu
	210					215					220		•	Ser	
225		•			230					235		-		Cys	240
Ser	Thr	Leu	Glu	Va1 245	Gln	Thr	Ser	Ala	Trp 250		Ala	Lys	Val	Ile 255	Phe
,	·		260					265					270	Ala	
		275		_			280			_		285		His	
Ala	Asn 290	Leu	Val	Leu	Ile	Tyr 295		Asn	Glu	Ile	Gly 300	Thr	Arg	Val	Phe
G1u 305		Thr	Glu	Thr	Ile 310	Thr	Leu	Asn	Phe	Ile 315		Phe	Ser	Met	Leu 320
•	·		_	325				·	330					Lys 335	
Ser	Asp	Asp	I1e 340	Ser	Val	Asn	Asp	Pro 345	Glu	His	Asn	Glu	A1a 350	-	Glu
Pro	His	Trp 355		Glu	Val	Glu	Gly 360	Gln	His	Leu	Gly	Cys 365		Ser	His
Leu	Met 370	Asp	Ala	Val	Cys	Gly 375		Glu	Gln	Arg	Asp 380		Asp	Thr	Ser

WO 01/46261 PCT/US00/35305

Leu Thr Gln His Gly Trp Leu Asn Ser Thr Ile Pro Thr Gly Glu Thr 390 395 Asp Thr Glu Pro Gln Tyr Lys Val Leu Ser Asp Phe Tyr Gly Glu Gly 405 410 Glu Ile Gln Leu Ser Cys Glu Pro Glu Glu Ala Ala Arg Thr Glu Lys 420 425 Ile Ser Glu Pro Leu Val Thr Ser Ala Asn Leu Asp Pro Gln Leu Glu 440 Asp Leu His His Leu Gly Gln Glu His Thr Val Ser Glu Asp Gly Pro 455 460 Glu Glu Glu Thr Ser Ile Thr Val Val Asp Trp Asp Pro Gln Thr Gly 470 475 Arg Leu Cys Ile Pro Ser Leu Pro Ile Phe Gly Arg Asp Pro Glu Asn 485 490 Tyr Gly His Tyr Glu Arg Asp Gln Leu Leu Glu Gly Gly Leu Leu Ser 505 Arg Leu Tyr Glu Asn Gln Ala Pro Asp Lys Pro Glu Lys Glu Asn Glu 520 525 Asn Cys Leu Thr Arg Phe Met Glu Glu Trp Gly Leu His Val Gln Met 530 535 540 Glu Ser 545 <210> 38 <211> 217 <212> PRT <213> Mus musculus <400> 38 Val Pro Cys Val Phe Cys Gly Leu Pro Lys Pro Thr Asn Ile Thr Phe Leu Ser Ile Asn Met Lys Asn Val Leu His Trp Asn Pro Pro Glu Ser 20 25 Leu His Gly Val Glu Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly Gln Lys Lys Trp Leu Asn Ala Ser Lys Cys Gly Ser Ile Asn Arg Thr Tyr Cys Asp Leu Ser Val Glu Thr Ser Asp Tyr Glu His Gln Phe Tyr 75 70 Ala Lys Val Lys Ala Ile Trp Glu Ala Arg Cys Ser Glu Trp Ala Glu 90 Thr Glu Arg Phe Tyr Pro Phe Leu Glu Thr Gln Val Ser Pro Pro Glu 100 105 110

Ile Ala Leu Thr Thr Gly Glu Lys Ser Ile Ser Ile Ala Leu Thr Ala 120 Pro Glu Lys Trp Lys Arg Asn Pro Gln Asp His Thr Val Ser Met Gln 135 140 Gln Ile Tyr Pro Asn Leu Lys Tyr Asn Val Ser Val Tyr Asn Thr Lys 155 Ser Arg Arg Thr Trp Ser Gln Cys Val Thr Asn Ser Thr Leu Val Leu 170 165 Ser Trp Leu Glu Pro Asn Thr Leu Tyr Cys Val His Val Glu Ser Leu 185 Val Pro Gly Pro Pro Arg Leu Pro Met Pro Ser Gln Lys Gln Cys Ile 200 205 Ser Thr Leu Glu Val Gln Thr Ser Ala 210 215 <210> 39 <211> 514

<212> PRT

<213> Mus musculus

<400> 39

Val Pro Cys Val Phe Cys Gly Leu Pro Lys Pro Thr Asn Ile Thr Phe 5 Leu Ser Ile Asn Met Lys Asn Val Leu His Trp Asn Pro Pro Glu Ser Leu His Gly Val Glu Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly 45 35 40 Gln Lys Lys Trp Leu Asn Ala Ser Lys Cys Gly Ser Ile Asn Arg Thr 55 60 Tyr Cys Asp Leu Ser Val Glu Thr Ser Asp Tyr Glu His Gln Phe Tyr 75 Ala Lys Val Lys Ala Ile Trp Glu Ala Arg Cys Ser Glu Trp Ala Glu Thr Glu Arg Phe Tyr Pro Phe Leu Glu Thr Gln Val Ser Pro Pro Glu 100 105 Ile Ala Leu Thr Thr Gly Glu Lys Ser Ile Ser Ile Ala Leu Thr Ala 120 125 Pro Glu Lys Trp Lys Arg Asn Pro Gln Asp His Thr Val Ser Met Gln 130 140 135 Gln Ile Tyr Pro Asn Leu Lys Tyr Asn Val Ser Val Tyr Asn Thr Lys 150 155

Ser Arg Arg Thr Trp Ser Gln Cys Val Thr Asn Ser Thr Leu Val Leu

170

175

WO 01/46261 PCT/US00/35305

			180)				185	5				190)	Leu
		195)				200)				205	G1r	Cys	Ile
	210					215					220	Lys	Val		Phe
225					230					235					Ile 240
				245					250					255	Pro
			260					265					270		Phe
		275					280			Ile		285			
	290					295				Asn	300				
305					310					His 315					320
				325					330	Leu				335	
			340					345		Arg			350		
		355					360			Ile Asp		365			
	370					375				ASP	380				
385					390					395 Leu					400
				405					410	Val				415	
			420					425		Trp			430		
Arg		435					440					445			
Tyr	450					455					460				
465 Arg					470					475					480
Asn				485					490					495	
Glu :			500					505	•	·			510		

<pre><210> 40 <211> 18 <212> DNA <213> Mus musculus</pre>	
<400> 40 egeogegtte cegagatg	18
<210> 41 <211> 24 <212> DNA <213> Mus musculus	
<400> 41 ggatgaggca gggctgacaa agtt	24
<210> 42 <211> 36 <212> DNA <213> Homo sapiens	
<400> 42 acttgtggaa ttcgctagca ccaagggccc atcggt	36
<210> 43 <211> 32 <212> DNA <213> Homo sapiens	
<400> 43 gcctagaacg cgttcattta cccggagaca gg	32
<210> 44 <211> 8 <212> DNA <213> Homo sapiens	
<400> 44 aattgaga	8
<210> 45	

	> DNA > Homo sapiens	
<400: cgcgtctc	> 45 ′	8
<400		37
<400 gacctgaacg	> 47 cgtctaacac tctcccctgt tg	32
<400: tcagtcggaa	> 48 ttcgcagaag ccatgcgggc tcccggcc	38
<400: ctgtgacgct	> 49 agcctctgat gattgatctt tcaaa	35

gato		100> gaa t	50 ctcgc	agaa	ag co	atgo	agac	ttt	caca	atg	gtt					43
	<2 <2	210> 211> 212> 213>	86	sap	oiens	;										
	acggt							a gat	tggcg	gga	agat	gaaq	jac a	agato	ggtgca	ı 60 86
	<2 <2	212>	1720		oiens	5										
	<2	220> 221> 222>	CDS (1).	(:	1713))										
	cgg		52 ccc Pro													48
			cct Pro 20				_	-		_	_			Gly	_	96
			gca Ala													144
			act Thr								_		-			192
			tat Tyr													240
gaa	tgc	aga	aat	atc	aat	aga	acc	tac	tgt	gat	ctt	tct	gct	gaa	act	288

Glu	Cys	Arg	Asn	Ile 85	Asn	Arg	Thr	Tyr	Cys 90	Asp	Leu	Ser	Ala	Glu 95	Thr	
	_		_		-			-	aaa Lys							336
									gga Gly							384
_									gca Ala							432
			-	-	_		_		gag Glu			-	-			480
	-								ata Ile 170							528
									aac Asn							576
-									tgg Trp							624
	_	Val							cca Pro							672
-									act Thr							720
	_	_			Gly		_	-	ttc Phe 250	Pro					Ser	768

							ggc Gly						816
							aac Asn						864
							cag Gln						912
			Thr	-			agc Ser 315	_	_			-	960
							agc Ser						1008
							act Thr					_	1056
							tca Ser						1104
	Pro	 _	Leu	-	Ile	Ser	cgg Arg			-	_		1152
		 -	 -		-	_	cct Pro 395		_	_			1200
	_						gcc Ala	-		_	_		1248
							gtc Val					-	1296

ctg cac cag gac tgg ctg aat ggc aag gag tac aag tgc aag gtc tcc Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser 435 440 445	1344
aac aaa gcc ctc cca gcc ccc atc gag aaa acc atc tcc aaa gcc aaa Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys 450 455 460	1392
ggg cag ccc cga gaa cca cag gtg tac acc ctg ccc cca tcc cgg gat Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp 465 470 475 480	1440
gag ctg acc aag aac cag gtc agc ctg acc tgc ctg gtc aaa ggc ttc Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe 485 490 495	1488
tat ccc agc gac atc gcc gtg gag tgg gag agc aat ggg cag ccg gag Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu 500 505 510	1536
aac aac tac aag acc acg cct ccc gtg ctg gac tcc gac ggc tcc ttc Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe 515 520 525	1584
ttc ctc tac agc aag ctc acc gtg gac aag agc agg tgg cag cag ggg Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly 530 535 540	1632
aac gtc ttc tca tgc tcc gtg atg cat gag gct ctg cac aac cac tac Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr 545 550 555 560	1680
acg cag aag agc ctc tcc ctg tct ccg ggt aaa tgacgcg Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys 565 570	1720
<210> 53 <211> 571 <212> PRT <213> Homo sapiens	
<400> 53	

PCT/US00/35305

Met 1	Arg	Ala	Pro	Gly 5	Arg	Pro	Ala	Leu	Arg 10	Pro	Leu	Leu	Leu	Leu 15	Leu
Leu	Ala	Ala	Pro 20	Trp	Gly	Arg	Ala	Val 25	Pro	Cys	Val	Ser	Gly 30	Gly	Leu
Pro	Lys	Pro 35	Ala	Asn	Ile	Thr	Phe 40	Leu	Ser	Ile	Asn	Met 45	Lys	Asn	Val
Leu	G1n 50	Trp	Thr	Pro	Pro	G1u 55	Gly	Leu	Gln	Gly	Va1 60	Lys	Val	Thr	Tyr
Thr 65	Val	Gln	Tyr	Phe	Ile 70	Tyr	Gly	Gln	Lys	Lys 75	Trp	Leu	Asn	Lys	Ser 80
Glu	Cys	Arg	Asn	Ile 85	Asn	Arg	Thr	Tyr	Cys 90	Asp	Leu	Ser	Ala	G1u 95	Thr
Ser	Asp	Tyr	Glu 100	His	Gln	Tyr	Tyr	Ala 105	Lys	Val	Lys	Ala	Ile 110	Trp	Gly
Thr	Lys	Cys 115	Ser	Lys	Trp	Ala	Glu 120	Ser	G1y	Arg	Phe	Tyr 125	Pro	Phe	Leu
Glu	Thr 130	Gln	Ile	Gly	Pro	Pro 135	Glu	Val	Ala	Leu	Thr 140	Thr	Asp	G1u	Lys
Ser 145	Ile	Ser	Val	Val	Leu 150	Thr	A1 a	Pro	Glu	Lys 155	Trp	Lys	Arg	Asn	Pro 160
Glu	Asp	Leu	Pro	Val 165	Ser	Met	Gln	Gln	Ile 170	Tyr	Ser	Asn	Leu	Lys 175	Tyr
Asn	Val	Ser	Val 180	Leu	Asn	Thr	Lys	Ser 185	Asn	Arg	Thr	Trp	Ser 190	Gln	Cys
Val	Thr	Asn 195	His	Thr	Leu	Val	Leu 200	Thr	Trp	Leu	Glu	Pro 205	Asn	Thr	Leu
Tyr	Cys 210	Val	His	Val	Glu	Ser 215	Phe	Val	Pro	Gly	Pro 220	Pro	Arg	Arg	Ala
G1n 225	Pro	Ser	Glu	Lys	G1n 230	Cys	Ala	Arg	Thr	Leu 235	Lys	Asp	Gln	Ser	Ser 240
Glu	Ala	Ser	Thr	Lys 245	Gly	Pro	Ser	Val	Phe 250	Pro	Leu	Ala	Pro	Ser 255	Ser
Lys	Ser	Thr	Ser 260	Gly	Gly	Thr	Ala	A1a 265	Leu	Gly	Cys	Leu	Va1 270	Lys	Asp
Tyr	Phe	Pro 275	Glu	Pro	Val	Thr	Va1 280	Ser	Trp	Asn	Ser	G1y 285	Ala	Leu	Thr
Ser	G1y 290	Val	His	Thr	Phe	Pro 295	Ala	Val	Leu	Gln	Ser 300	Ser	Gly	Leu	Tyr
Ser 305	Leu	Ser	Ser	Val	Val 310	Thr	Val	Pro	Ser	Ser 315	Ser	Leu	G1y	Thr	G1n 320
Thr	Tyr	Ile	Cys	Asn 325	Val	Asn	His	Lys	Pro 330	Ser	Asn	Thr	Lys	Val 335	Asp

WO 01/46261 PCT/US00/35305

350 345 340 Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro 365 Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr 375 380 Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn 395 390 385 Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg 405 410 Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val 425 430 Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser 440 Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys 460 455 Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp 470 475 Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe 490 Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu 505 Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe 525 515 520 Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly 540 535 Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr 560 555 550 Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys 565 570 <210> 54

<211> 547

<212> PRT

<213> Homo sapiens

<400> 54

Val Pro Cys Val Ser Gly Gly Leu Pro Lys Pro Ala Asn Ile Thr Phe 1 5 10 15 Leu Ser Ile Asn Met Lys Asn Val Leu Gln Trp Thr Pro Pro Glu Gly 20 25 30 Leu Gln Gly Val Lys Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly 35 40 45

WO 01/46261 PCT/US00/35305

Gln	Lys 50	Lys	Trp	Leu	Asn	Lys 55	Ser	Glu	Cys	Arg	Asn 60	Ile	Asn	Arg	Thr
Tyr 65	Cys	Asp	Leu	Ser	Ala 70	Glu	Thr	Ser	Asp	Tyr 75	Glu	His	Gln	Tyr	Tyr 80
Ala	Lys	Val	Lys	A1 a 85	Ile	Trp	Gly	Thr	Lys 90	Cys	Ser	Lys	Trp	A1a 95	Glu
Ser	Gly	Arg	Phe 100	Tyr	Pro	Phe	Leu	Glu 105	Thr	Gln	Ile	Gly	Pro 110	Pro	Glu
۷a۱	Ala	Leu 115	Thr	Thr	Asp	Glu	Lys 120	Ser		Ser	Val	Val 125	Leu	Thr	Ala
Pro	Glu 130		Trp	Lys	Arg	Asn 135	Pro			Leu	Pro 140	Val	Ser	Met	Gln
G1n 145		Tyr	Ser	Asn	Leu 150	Lys	Tyr	Asn	Val	Ser 155	Val	Leu	Asn	Thr	Lys 160
	Asn	Arg	Thr	Trp 165	Ser	Gln	Cys	Val	Thr 170	Asn		Thr	Leu	Val 175	Leu
Thr	Trp	Leu	G1u 180		Asn	Thr	Leu	Tyr 185		Val	His	Val	Glu 190	Ser	Phe
Val	Pro	Gly 195	Pro		Arg	Arg	Ala 200			Ser	Glu	Lys 205	Gln	Cys	Ala
Arg	Thr 210		Lys	Asp	Gln	Ser 215		Glu	Ala	Ser	Thr 220	Lys	Gly	Pro	Ser
Va1 225		Pro	Leu	Ala	Pro 230		Ser	Lys	Ser	Thr 235		Gly	Gly	Thr	A1a 240
Ala	Leu	Gly	Cys	Leu 245		Lys	Asp	Tyr	Phe 250			Pro	Val	Thr 255	Val
Ser	Trp	Asn	Ser 260		Ala	Leu	Thr	Ser 265		Val	His	Thr	Phe 270	Pro	Ala
Val	Leu	G1n 275		Ser	Gly	Leu	Tyr 280		Leu	Ser	Ser	Va1 285		Thr	Val
Pro	Ser 290		Ser	Leu	Gly	Thr 295		Thr	Tyr	· Ile	Cys 300		Val	Asn	His
Lys 305		Ser	Asn	Thr	Lys 310		Asp	Lys	Lys	Val 315		Pro	Lys	Ser	Cys 320
Asp	Lys	Thr	His	Thr 325		Pro	Pro	Cys	Pro 330		Pro	Glu	ı Leu	Leu 335	
Gly	Pro	Ser	Val 340		. Leu	Phe	Pro	Pro 345		Pro	Lys	Asp	Thr 350	Leu	Met
Πe	Ser	Arg 355		Pro	Glu	\Val	Thr 360		: Val	۷a٦	۷a۱	Asp 365		Ser	His
G٦ι	Asp 370	Pro		ı Val	Lys	275 375		ı Trp	Tyr	· Val	Asp 380		√ Val	Glu	Va1

PCT/US00/35305 WO 01/46261

His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr 390 395 Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly 405 410 Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile 425 Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val 440 Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser 450 455 460 Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu 475 Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro 485 490 Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val 505 Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met 520 His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser 530 535 Pro Gly Lys 545 <210> 55 <211> 217 <212> PRT <213> Homo sapiens <400> 55 Val Pro Cys Val Ser Gly Gly Leu Pro Lys Pro Ala Asn Ile Thr Phe Leu Ser Ile Asn Met Lys Asn Val Leu Gln Trp Thr Pro Pro Glu Gly Leu Gln Gly Val Lys Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly Gln Lys Lys Trp Leu Asn Lys Ser Glu Cys Arg Asn Ile Asn Arg Thr Tyr Cys Asp Leu Ser Ala Glu Thr Ser Asp Tyr Glu His Gln Tyr Tyr 70 75 Ala Lys Val Lys Ala Ile Trp Gly Thr Lys Cys Ser Lys Trp Ala Glu 90 Ser Gly Arg Phe Tyr Pro Phe Leu Glu Thr Gln Ile Gly Pro Pro Glu 100 105 110

Val	Ala	Leu 115	Thr	Thr	Asp	Glu	Lys 120	Ser	Ile	Ser	Val	Val 125	Leu	Thr	Ala	
Pro	Glu 130		Trp	Lys	Arg	Asn 135		G1u	Asp	Leu	Pro 140		Ser	Met	Gln	
G1n 145	Ile	Tyr	Ser	Asn	Leu 150	Lys	Tyr	Asn	Val	Ser 155		Leu	Asn	Thr	Lys 160	
Ser	Asn	Arg	Thr	Trp 165	Ser	Gln	Cys	Val	Thr 170	Asn	His	Thr	Leu	Val 175	Leu	
Thr	Trp	Leu	Glu 180	Pro	Asn	Thr	Leu	Tyr 185	Cys	Val	His	Val	Glu 190	Ser	Phe	
Val	Pro	Gly 195	Pro	Pro	Arg	Arg	Ala 200	Gln	Pro	Ser	Glu	Lys 205	Gln	Cys	Ala	
Arg	Thr 210	Leu	Lys	Asp	Gln	Ser 215	Ser	Glu								
	<2 <2	212>	101: DNA	l o sap	oiens	5										
	<2	220> 221> 222>		(:	1008)										
2+4		100>					_4_				.			-4.1	1.1	40
													agt Ser			48
													gat Asp 30			96
													acc Thr			144
													gaa Glu			192
tac	tat	tct	gtc	gaa	tac	cag	ggġ	gag	tac	gag	agc	ctg	tac	acg	agc	240

Tyr 65	Tyr	Ser	Val	Glu	Tyr 70	Gln	Gly	Glu	Tyr	G1u 75	Ser	Leu	Tyr	Thr	Ser 80		
				ccc Pro 85												2	288
				gat Asp												(336
				ttg Leu			_			_		_		_	•	;	384
				aga Arg												4	432
				ggc Gly												4	480
				ttc Phe 165													528
				aaa Lys													576
				cca Pro							_	-	_			(624
				999 Gly			_	_		-	_		_	-	-	(672
				gag Glu													720

WO 01/46261 53

ccg cca tct Pro Pro Ser					_		•		-	768
ctg ctg aat Leu Leu Asn						-		-		816
gat aac gcc Asp Asn Ala 275		Ser Gly								864
gac agc aag Asp Ser Lys 290			-	_	_	Leu	_	-	-	912
aaa gca gac Lys Ala Asp 305	Tyr Glu l		_	Tyr A		_	-			960
cag ggc ctg Gln Gly Leu	-	_	-	-					-	1008
tag										1011
<210> <211> <212> <213>	336	iens								
<400>										
Met Gln Thr 1	Phe Thr N	Met Val	Leu Glu	Glu :	Ile Trp) Thr	Ser	Leu 15	Phe	
Met Trp Phe	Phe Tyr A	Ala Leu	Ile Pro 25	Cys I	Leu Lei	ı Thr	Asp 30	Glu	Val	
Ala Ile Leu 35		Pro Gln		Ser \	Val Lei	J Ser 45		Asn	Met	
Lys His Leu 50	Leu Met ⁻	Trp Ser 55		Ile /	Ala Pro 60		G1u	Thr	Val	
Tyr Tyr Ser 65			Gly Glu	-		r Leu	Tyr	Thr	Ser 80	

His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr Asn Leu Arg 105 Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser Ile Leu Lys 120 His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro Gly Met Glu 135 Ile Thr Lys Asp Gly Phe His Leu Val Ile Glu Leu Glu Asp Leu Gly 150 155 Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Arg Glu Pro Gly Ala 170 Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro Val His Leu 185 Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala Gln Thr Phe 200 Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr Glu Cys Val 215 220 Glu Val Gln Gly Glu Ala Thr Val Ala Ala Pro Ser Val Phe Ile Phe 230 235 Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys 250 Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val 265 Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln 275 280 285 Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser 295 300 Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His 310 315 Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys 325 330 <210> 58

<211> 307

<212> PRT

<213> Homo sapiens

<400> 58

Asp Glu Val Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser 1 5 10 15 Thr Asn Met Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly 20 25 30

```
Glu Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu
                            40
Tyr Thr Ser His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu
Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr
                    70
                                        75
Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser
                                    90
Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro
            100
                                105
Gly Met Glu Ile Thr Lys Asp Gly Phe His Leu Val Ile Glu Leu Glu
                            120
Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Arg Arg Glu
                        135
                                            140
Pro Gly Ala Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro
                    150
                                        155
Val His Leu Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala
                165
                                    170
Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr
            180
Glu Cys Val Glu Val Gln Gly Glu Ala Thr Val Ala Ala Pro Ser Val
                            200
                                                205
Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser
                        215
                                            220
Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln
225
                    230
                                        235
Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val
                245
                                    250
Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu
            260
                                265
                                                     270
Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu
        275
                            280
                                                285
Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg
    290
                        295
                                            300
Gly Glu Cys
305
```

<210> 59

<211> 201

<212> PRT

<213> Homo sapiens

<400> 59

Asp Glu Val Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser Thr Asn Met Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly Glu Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu Tyr Thr Ser His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu 55 Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro 105 110 Gly Met Glu Ile Thr Lys Asp Gly Phe His Leu Val Ile Glu Leu Glu 125 120 Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Arg Arg Glu 135 140 Pro Gly Ala Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro 150 160 145 Val His Leu Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala 170 165 Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr 190 180 185 Glu Cys Val Glu Val Gln Gly Glu Ala 195 200 <210> 60 <211> 323 <212> PRT <213> Homo sapiens <400> 60 Asp Glu Val Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser 10 Thr Asn Met Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly Glu Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu 45 40 Tyr Thr Ser His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu 55

Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr

70

Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro 100 105 Gly Met Glu Ile Pro Lys His Gly Phe His Leu Val Ile Glu Leu Glu 120 Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Thr Arg Glu 135 140 Pro Gly Ala Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro 150 155 Val His Leu Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala 165 170 Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr 180 185 Glu Cys Val Glu Val Gln Gly Glu Ala Gly Gly Gly Gly Ser Gly Gly 200 Gly Gly Ser Gly Gly Gly Ser Arg Thr Val Ala Ala Pro Ser Val 215 Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser 230 Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln 245 250 Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val 260 265 Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu 275 280 285 Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu 295 Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg 305 310 315 320 Gly Glu Cys <210> 61 <211> 201 <212> PRT <213> Homo sapiens <400> 61 Asp Glu Val Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser 10

Thr Asn Met Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly

25

30

PCT/US00/35305 WO 01/46261

Glu Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu Tyr Thr Ser His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr 75 Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser 85 90 Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro 105 Gly Met Glu Ile Pro Lys His Gly Phe His Leu Val Ile Glu Leu Glu 120 125 Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Thr Arg Glu 135 130 Pro Gly Ala Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro 150 155 Val His Leu Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala 165 170 Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr 190 180 Glu Cys Val Glu Val Gln Gly Glu Ala 195 200 <210> 62 <211> 559 <212> PRT <213> Homo sapiens <400> 62 Val Pro Cys Val Ser Gly Gly Leu Pro Lys Pro Ala Asn Ile Thr Phe Leu Ser Ile Asn Met Lys Asn Val Leu Gln Trp Thr Pro Pro Glu Gly 20 Leu Gln Gly Val Lys Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly 40 Gln Lys Lys Trp Leu Asn Lys Ser Glu Cys Arg Asn Ile Asn Arg Thr 60 Tyr Cys Asp Leu Ser Ala Glu Thr Ser Asp Tyr Glu His Gln Tyr Tyr 70 65 75 Ala Lys Val Lys Ala Ile Trp Gly Thr Lys Cys Ser Lys Trp Ala Glu 90 85

Ser Gly Arg Phe Tyr Pro Phe Leu Glu Thr Gln Ile Gly Pro Pro Glu

105

110

Val	Ala	Leu 115	Thr	Thr	Asp	Glu	Lys 120	Ser	Ile	Ser	Val	Va I 125	Leu	Ihr	Ala
Pro	Glu 130	Lys	Trp	Lys	Arg	Asn 135	Pro	Glu	Asp	Leu	Pro 140	Val	Ser	Met	Gln
145	Пе			Asn	150					155					160
				Trp 165					170					175	
	•		180	Pro				185					190		
۷a٦	Pro	Gly 195	Pro	Pro	Arg	Arg	Ala 200	Gln	Pro	Ser	Glu	Lys 205	G1n	Cys	Ala
	210			Asp		215					220				
225				Ser	230					235					240
				Lys 245					250					255	
			260					265					270		Ser
		275					280					285			Ser
	290					295	+				300				Ser
305)				310					315)				Asn 320
				325)				330)				335	
	_		340)				345	·)				350)	Val
		355	5				360)				365	<u>, </u>		Thr
	370)				375	5				380)			Glu
389	5				390)				395	5				400
				409	5				410)				415	
			420)				425	5				430)	Lys
Cy	s Ly:	s Va ⁻ 43!		r Ası	n Lys	s Ali	a Lei 44(s Se	r Sei	r Ile	e Glu 44!	u Lys 5	s Thi	^ Ile

Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu 465 470 475 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn 490 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser 505 500 510 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg 520 525 Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys 545 550

<210> 63 <211> 214

<212> PRT

<213> Homo sapiens

165

<400> 63

Val Pro Cys Val Ser Gly Gly Leu Pro Lys Pro Ala Asn Ile Thr Phe Leu Ser Ile Asn Met Lys Asn Val Leu Gln Trp Thr Pro Pro Glu Gly Leu Gln Gly Val Lys Val Thr Tyr Thr Val Gln Tyr Phe Ile Tyr Gly 40 Gln Lys Lys Trp Leu Asn Lys Ser Glu Cys Arg Asn Ile Asn Arg Thr Tyr Cys Asp Leu Ser Ala Glu Thr Ser Asp Tyr Glu His Gln Tyr Tyr Ala Lys Val Lys Ala Ile Trp Gly Thr Lys Cys Ser Lys Trp Ala Glu 90 Ser Gly Arg Phe Tyr Pro Phe Leu Glu Thr Gln Ile Gly Pro Pro Glu 105 100 Val Ala Leu Thr Thr Asp Glu Lys Ser Ile Ser Val Val Leu Thr Ala 120 125 Pro Glu Lys Trp Lys Arg Asn Pro Glu Asp Leu Pro Val Ser Met Gln 130 135 140 Gln Ile Tyr Ser Asn Leu Lys Tyr Asn Val Ser Val Leu Asn Thr Lys 150 155 Ser Asn Arg Thr Trp Ser Gln Cys Val Thr Asn His Thr Leu Val Leu

170

```
Thr Trp Leu Glu Pro Asn Thr Leu Tyr Cys Val His Val Glu Ser Phe
                                185
Val Pro Gly Pro Pro Arg Arg Ala Gln Pro Ser Glu Lys Gln Cys Ala
        195
                            200
                                                205
Arg Thr Leu Lys Asp Gln
    210
      <210> 64
      <211> 19
      <212> PRT
      <213> Homo sapiens
      <400> 64
Glu Glu Ile His Ala Glu Leu Arg Arg Phe Arg Arg Val Pro Cys Val
1
                                    10
Ser Gly Gly
      <210> 65
      <211> 207
      <212> PRT
      <213> Homo sapiens
      <400> 65
Leu Pro Lys Pro Ala Asn Ile Thr Phe Leu Ser Ile Asn Met Lys Asn
Val Leu Gln Trp Thr Pro Pro Glu Gly Leu Gln Gly Val Lys Val Thr
                                25
Tyr Thr Val Gln Tyr Phe Ile Tyr Gly Gln Lys Lys Trp Leu Asn Lys
                            40
Ser Glu Cys Arg Asn Ile Asn Arg Thr Tyr Cys Asp Leu Ser Ala Glu
                        55
Thr Ser Asp Tyr Glu His Gln Tyr Tyr Ala Lys Val Lys Ala Ile Trp
65
                    70
                                        75
Gly Thr Lys Cys Ser Lys Trp Ala Glu Ser Gly Arg Phe Tyr Pro Phe
Leu Glu Thr Gln Ile Gly Pro Pro Glu Val Ala Leu Thr Thr Asp Glu
                                105
Lys Ser Ile Ser Val Val Leu Thr Ala Pro Glu Lys Trp Lys Arg Asn
                                                 125
                            120
Pro Glu Asp Leu Pro Val Ser Met Gln Gln Ile Tyr Ser Asn Leu Lys
                        135
                                             140
Tyr Asn Val Ser Val Leu Asn Thr Lys Ser Asn Arg Thr Trp Ser Gln
145
                    150
                                                             160
```

```
Cys Val Thr Asn His Thr Leu Val Leu Thr Trp Leu Glu Pro Asn Thr
                165
                                    170
Leu Tyr Cys Val His Val Glu Ser Phe Val Pro Gly Pro Pro Arg Arg
            180
                                185
Ala Gln Pro Ser Glu Lys Gln Cys Ala Arg Thr Leu Lys Asp Gln
                            200
      <210> 66
      <211> 150
      <212> PRT
      <213> Homo sapiens
      <400> 66
Cys Arg Asn Ile Asn Arg Thr Tyr Cys Asp Leu Ser Ala Glu Thr Ser
Asp Tyr Glu His Gln Tyr Tyr Ala Lys Val Lys Ala Ile Trp Gly Thr
Lys Cys Ser Lys Trp Ala Glu Ser Gly Arg Phe Tyr Pro Phe Leu Glu
                            40
Thr Gln Ile Gly Pro Pro Glu Val Ala Leu Thr Thr Asp Glu Lys Ser
                        55
Ile Ser Val Val Leu Thr Ala Pro Glu Lys Trp Lys Arg Asn Pro Glu
Asp Leu Pro Val Ser Met Gln Gln Ile Tyr Ser Asn Leu Lys Tyr Asn
                                     90
Val Ser Val Leu Asn Thr Lys Ser Asn Arg Thr Trp Ser Gln Cys Val
            100
                                 105
Thr Asn His Thr Leu Val Leu Thr Trp Leu Glu Pro Asn Thr Leu Tyr
                            120
Cys Val His Val Glu Ser Phe Val Pro Gly Pro Pro Arg Arg Ala Gln
                        135
                                             140
Pro Ser Glu Lys Gln Cys
145
                     150
      <210> 67
      <211> 196
      <212> PRT
      <213> Homo sapiens
      <400> 67
Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser Thr Asn Met Lys His
                                                         15
                                     10
```

Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly Glu Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu Tyr Thr Ser His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr Asn Leu Arg Val Arg 75 70 Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro Gly Met Glu Ile Thr 100 105 Lys Asp Gly Phe His Leu Val Ile Glu Leu Glu Asp Leu Gly Pro Gln 120 115 Phe Glu Phe Leu Val Ala Tyr Trp Arg Arg Glu Pro Gly Ala Glu Glu 135 His Val Lys Met Val Arg Ser Gly Gly Ile Pro Val His Leu Glu Thr 150 155 Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr Glu Cys Val Glu Val 185 190 Gln Gly Glu Ala 195 <210> 68 <211> 203 <212> PRT <213> Homo sapiens <400> 68 Asp Glu Val Ala Ile Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser Thr Asn Met Lys His Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly 25 Glu Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu 40 Tyr Thr Ser His Ile Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu 55 Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr 75 70 Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser

90

85

Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro 105 Gly Met Glu Ile Pro Lys His Gly Phe His Leu Val Ile Glu Leu Glu 115 120 125 Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala Tyr Trp Thr Arg Glu 135 Pro Gly Ala Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro 155 150 Val His Leu Glu Thr Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala 170 Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr 180 185 Glu Cys Val Glu Val Gln Gly Glu Ala Ile Pro 195 200 <210> 69 <211> 196 <212> PRT <213> Homo sapiens <400> 69 Leu Pro Ala Pro Gln Asn Leu Ser Val Leu Ser Thr Asn Met Lys His 5 Leu Leu Met Trp Ser Pro Val Ile Ala Pro Gly Glu Thr Val Tyr Tyr Ser Val Glu Tyr Gln Gly Glu Tyr Glu Ser Leu Tyr Thr Ser His Ile 40 Trp Ile Pro Ser Ser Trp Cys Ser Leu Thr Glu Gly Pro Glu Cys Asp 55 Val Thr Asp Asp Ile Thr Ala Thr Val Pro Tyr Asn Leu Arg Val Arg 75 Ala Thr Leu Gly Ser Gln Thr Ser Ala Trp Ser Ile Leu Lys His Pro 90 Phe Asn Arg Asn Ser Thr Ile Leu Thr Arg Pro Gly Met Glu Ile Pro 105 Lys His Gly Phe His Leu Val Ile Glu Leu Glu Asp Leu Gly Pro Gln 120 125 Phe Glu Phe Leu Val Ala Tyr Trp Thr Arg Glu Pro Gly Ala Glu Glu 130 135 140 His Val Lys Met Val Arg Ser Gly Gly Ile Pro Val His Leu Glu Thr 150 155 Met Glu Pro Gly Ala Ala Tyr Cys Val Lys Ala Gln Thr Phe Val Lys

170

175

```
Ala Ile Gly Arg Tyr Ser Ala Phe Ser Gln Thr Glu Cys Val Glu Val
                                185
            180
Gln Gly Glu Ala
        195
      <210> 70
      <211> 135
      <212> PRT
      <213> Homo sapiens
      <400> 70
Cys Ser Leu Thr Glu Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr
Ala Thr Val Pro Tyr Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln
                                 25
Thr Ser Ala Trp Ser Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr
                            40
Ile Leu Thr Arg Pro Gly Met Glu Ile Thr Lys Asp Gly Phe His Leu
                        55
                                             60
Val Ile Glu Leu Glu Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala
                    70
                                         75
Tyr Trp Arg Arg Glu Pro Gly Ala Glu Glu His Val Lys Met Val Arg
Ser Gly Gly Ile Pro Val His Leu Glu Thr Met Glu Pro Gly Ala Ala
                                 105
Tyr Cys Val Lys Ala Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser
                             120
        115
                                                 125
Ala Phe Ser Gln Thr Glu Cys
    130
      <210> 71
      <211> 135
      <212> PRT
      <213> Homo sapiens
      <400> 71
Cys Ser Leu Thr Glu Gly Pro Glu Cys Asp Val Thr Asp Asp Ile Thr
                                     10
Ala Thr Val Pro Tyr Asn Leu Arg Val Arg Ala Thr Leu Gly Ser Gln
                                 25
Thr Ser Ala Trp Ser Ile Leu Lys His Pro Phe Asn Arg Asn Ser Thr
        35
```

Ile Leu Thr Arg Pro Gly Met Glu Ile Pro Lys His Gly Phe His Leu 55 Val Ile Glu Leu Glu Asp Leu Gly Pro Gln Phe Glu Phe Leu Val Ala 65 70 75 80 Tyr Trp Thr Arg Glu Pro Gly Ala Glu Glu His Val Lys Met Val Arg Ser Gly Gly Ile Pro Val His Leu Glu Thr Met Glu Pro Gly Ala Ala 100 105 110 Tyr Cys Val Lys Ala Gln Thr Phe Val Lys Ala Ile Gly Arg Tyr Ser 120 125 Ala Phe Ser Gln Thr Glu Cys 130 135 <210> 72 <211> 15 <212> PRT <213> Homo sapiens <400> 72 Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Ser 5 10 15

Int anal Application No PCT/US 00/35305

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07K16/24 C07K16/28 C07K14/54 A61K39/395 C07K19/00 A61P11/06 //C07K14/715 A61P17/06 According to International Patent Classification (IPC) or to both national classification and IPC **B. FIELDS SEARCHED** Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE C. DOCUMENTS CONSIDERED TO BE RELEVANT Category ° Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. X WO 99 27103 A (ZYMOGENETICS INC) 1,2,7,8 3 June 1999 (1999-06-03) abstract page 2, line 10-36 page 4, line 13-17 page 5, line 5-17 page 19, line 11-33 page 37, line 11-33 page 41, line 19-24 Y page 43, line 1-26 3-6Further documents are listed in the continuation of box C. Patent family members are listed in annex. X Special categories of cited documents: "T" later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the "A" document defining the general state of the art which is not considered to be of particular relevance invention "E" earlier document but published on or after the international "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another involve an inventive step when the document is taken alone document of particular relevance; the claimed invention citation or other special reason (as specified) cannot be considered to involve an inventive step when the document is combined with one or more other such docudocument referring to an oral disclosure, use, exhibition or ments, such combination being obvious to a person skilled other means document published prior to the international filing date but later than the priority date claimed in the art. "&" document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report **71**. 05. 01 25 April 2001 Name and mailing address of the ISA Authorized officer European Patent Office, P.B. 5818 Patentiaan 2 NL – 2280 HV Rijswijk Tel. (+31–70) 340–2040, Tx. 31 651 epo nl, Fax: (+31–70) 340–3016 Montrone, M

Int Inal Application No PCT/US 00/35305

1,2 3-6
3-6
3-6
1,2
3-6
1-6
1-6

Inte lal Application No PCT/US 00/35305

	INTERNATIONAL SEARCH REPORT	PCT/US 00/	35305
C.(Continua	tion) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages		Relevant to claim No.
P,X	WO 00 42189 A (MURPHY MARIANNE; FLORENCE KIMBERLY (US); HUMAN GENOME SCIENCES INC) 20 July 2000 (2000-07-20) page 3, line 4-17 page 5, line 27 -page 6, line 31 page 7, line 6-36 page 8, line 1-3 page 9, line 8-12 page 30, line 32-37 page 51, line 19 -page 53, line 26 page 55, line 9-14 page 57, line 9,10 page 62, line 23 -page 64, line 30 page 115, line 5 -page 116, line 25		1,2,9,10

ational application No. PCT/US 00/35305

Box I	Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)
This Inter	national Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely: Although claims 1 to 10 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged
	effects of the compound/composition. Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Fule 6.4(a).
Box II	Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
THIS HILE	mational Searching Authority found multiple inventions in this international application, as follows:
1.	As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
з. 🗌	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4.	No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the Invention first mentioned in the claims; it is covered by claims Nos.:
Remark	on Protest The additional search fees were accompanied by the applicant's protest. No protest accompanied the payment of additional search fees.

Information on patent family members

Inti anal Application No
PCT/US 00/35305

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9927103	A	03-06-1999	AU BR CN EP NO	1607799 A 9814904 A 1282372 T 1032671 A 20002698 A	15-06-1999 03-10-2000 31-01-2001 06-09-2000 20-07-2000
WO 9903982	A	28-01-1999	AU AU EP EP WO	8404598 A 8571198 A 1027430 A 1012260 A 9903990 A	10-02-1999 10-02-1999 16-08-2000 28-06-2000 28-01-1999
US 5945511	A	31-08-1999	AU EP WO	6535098 A 1009825 A 9837193 A	09-09-1998 21-06-2000 27-08-1998
WO 0042189	A	20-07-2000	AU	2965600 A	01-08-2000

